

Original Research

Evaluation of the effectiveness of educational medical informatics tutorial on improving pharmacy students' knowledge and skills about the clinical problem-solving process

Ahmad R. Alsayed , Luai Hasoun , Heba A Khader , Abdullah Al-Dulaimi , Alaa AbuAwad , Iman Basheti , Mohammed Al Maqbali 

Received (first version): 03-Mar-2022

Accepted: 01-Apr-2022

Published online: 05-Apr-2022

Abstract

Objective: To investigate the effectiveness of an online tutorial and its impact on improving knowledge and skills of pharmacy students in the clinical problem-solving process that is necessary to implement pharmaceutical care. **Methods:** This is a prospective interventional study conducted during the COVID-19 pandemic restrictions using four novel templates. The first two levels of Kirkpatrick's Model (Reaction and Learning) were used. **Results:** 129 participants completed all of the online training parts. The findings indicated a significant improvement in the students' knowledge and skills. The participants achieved higher score following the tutorial than the baseline, with a statistically significant difference ($p < 0.001$). There was a significant improvement in the number of detected treatment-related problems. The majority of students were satisfied with the overall training process and stated a high evaluation score out of 10 (mean = 7.93 ± 1.42 , median = 8.00). **Conclusion:** The educational intervention achieved a substantial positive impact on decision-making skills of participating students and was considered effective in helping them attain basic skills such as teamwork, peer assessment, communication and critical evaluation. Healthcare providers must work together to ensure accurate medication use during care transitions. Pharmacists, as medication experts, play an important role in the implementation process. Pharmacy educators must prepare pharmacy student to use pharmaceutical care in their future practice.

Keywords: Clinical problem-solving process; Pharmacy; Informatics; Online learning

Ahmad R. ALSAYED*. PharmD, MSc, PhD. Department of Clinical Pharmacy and Therapeutics, Applied Science Private University, Amman 11931-166, Jordan. a_alsayed@asu.edu.jo

Luai HASOUN. PhD. Department of Clinical Pharmacy and Therapeutics, Applied Science Private University, Amman 11931-166, Jordan. l_hason@asu.edu.jo

Heba A KHADER. PhD. Department of Clinical Pharmacy and Pharmacy Practice, Faculty of Pharmaceutical Sciences, The Hashemite University, PO Box 330127, Zarqa 13133, Jordan. hebaa@hu.edu.jo

Abdullah AL-DULAIMI. MSc. Department of Clinical Pharmacy and Therapeutics, Applied Science Private University, Amman 11931-166, Jordan. abedhatem97@gmail.com

Alaa ABUAWAD. PhD. Department of Pharmaceutical Sciences and Pharmaceutics, Applied Science Private University, Amman 11931-166, Jordan. a_abuawad@asu.edu.jo

Iman BASHETI. PhD. Department of Clinical Pharmacy and Therapeutics, Applied Science Private University, Amman 11931-166, Jordan. dr_iman@asu.edu.jo

Mohammed AL MAQBALI. PhD. Department of Nursing Midwifery and Health, Northumbria University, Newcastle-Upon-Tyne, UK. mohammed.maqbali@northumbria.ac.uk

INTRODUCTION

Over years, the role of pharmacists has expanded from traditional dispensing to patient-focused pharmaceutical care (PC) practice.^{1,2} Pharmacists are now working collaboratively with other healthcare practitioners as a neighborhood of the health-care team to provide optimized patient care services in hospitals and various clinical settings.³⁻⁷ As pharmacists are now required to possess closer contact with patients,⁸ the general public naturally expects them to be able to handle challenges competently and in the best interests of their patients.⁹

The term PC was first defined by Hepler and Strand as "the responsible provision of drug therapy for the purpose of achieving definite outcomes that improve a patient's quality of life".¹⁰ More recently, PC was defined by the Pharmaceutical Care Network Europe as "the pharmacist's contribution to care of individuals in order to optimize medicines use and improve health outcomes".¹¹

Many studies have determined that involvement of pharmacists towards PC services improves patients' health outcomes through the identification and prevention of TRPs in several diseases¹²⁻¹⁹ and reduces health costs.²⁰⁻²³

However, several barriers have impeded the implementation of PC practice worldwide. These include lack of pharmacists' time, poor clinical knowledge and communication skills, insufficient pharmacists' self-confidence in addition to the negative attitudes of pharmacists themselves toward performing PC.²⁴⁻²⁹



International pharmacy bodies like The Accreditation Council for Pharmacy Education (ACPE)³⁰ and The Centre for the Advancement of Pharmacy Education (CAPE)³¹ require trainee (and practicing) pharmacists to possess an appropriate awareness and competencies within a patient-centred model of practice to enable evidence-based decision-making.^{31,32} This needs academically well-designed teaching components embedded within pharmacy curricula around evidence-based decision-making.

Accordingly, it's vital to organize pharmacy practitioners of today and tomorrow. Different studies have described the role that the educators of pharmacy school should play in supporting the PC practitioners.³³⁻³⁸ They have to design educational tools that promote the performance of pharmacists who are willing to implement PC in their practice and to ensure that students have a deep understanding of the principles and practices of PC.³⁹ In addition, they ought to establish in student's appropriate clinical knowledge, communication and problem-solving skills, and self-confidence to take the responsibility for providing PC.³⁹ One of the challenges faced by pharmacy educators is the motivation of students to provide PC.⁴⁰ However, enhancing students' tendency toward PC will be reflected on pharmacy educators readiness to increase PC's supply in practice.⁴⁰ This combination of knowledge, skills and attitudes is required for widespread acceptance and implementation of the concept of PC.

There is a recognized value of PC and clinical problem-solving as a process that reduces medication errors and patient harm.^{41,42} However, health care providers generally, pharmacists are not an exception, receive little formal education during their college years on how to collect patient's data, assess these data including the medications, design and implement the patient's care plan, educate the patient, with appropriate follow-up.^{41,42} This may be attributed to the lack of awareness and unclear understanding of who has the responsibility for providing this service.^{43,44} Therefore, this study aims to investigate the effectiveness of a tutorial training and its impact on improving knowledge and skills of pharmacy students in the clinical problem-solving process that is necessary to implement PC.

METHODS

Study setting and participants

This is a prospective interventional study that was conducted from October 2020 to February 2021 at the faculty of pharmacy at Applied Science Private University (ASU), a well-ranked Jordanian university located in Amman, Jordan. Pharmacy students enrolled in the Clinical Pharmacy and Therapeutics Lab module were invited to participate in the study during the study period.

This interventional trial was undertaken in the midst of the COVID-19 outbreak. As a result, researchers were forced to change the course and give it online via Microsoft Teams[®] rather than using the intended face-to-face delivery method.

The ethics committee of ASU granted approval under the

number 2021-PHA-4-5. All students were advised that participation was voluntary and that their responses would be kept completely confidential.

Designing and implementing the tutorial intervention

We developed the educational intervention in light of the available literature and the researchers' personal experiences. Prior to initiating the intervention, the material was reviewed and confirmed to ensure the information supplied was clear and accurate. Additionally, after developing the tutorial, it was pilot tested on 19 students during the preceding semester (not included in the analysis). Additionally, the material was re-evaluated following the pilot research.

The initial part of this study involved pre-testing students prior to delivering the educational intervention. The second part was the educational intervention's delivery. The third part involved administering the post-intervention test. Finally, the fourth part comprised a focus group discussion with the intervention's participants to get comments about the intervention. Three researchers (all with PhD, degree in clinical pharmacy and a minimum of seven years' experience in the clinical field) oversaw the educational intervention and data collection. Four innovative templates were used before, during, and following this instructional course: data collection (Appendix-Table A), basic calculations (Appendix -Table B), medication assessment (Appendix -Table C), and care plan (Appendix -Table D). Different versions of the tables in Appendix – Tables C and D were utilized depending on the patient's medication and disease status. These templates are part of a larger project coordinated by the first author that includes a website and a mobile application aimed at providing PC support (<https://www.asami-draacare.com>).

All surveys were distributed in English, the official language of pharmacy education in Jordanian universities. To ensure face validity, three academics with considerable experience conducting clinical studies and a broad spectrum of professional clinical experience examined the questionnaires prior to the current investigation.

Part 1: Pre- intervention part

After acquiring students' informed written agreement, a pre-training data collecting form and ability to analyze a clinical case was used to assess students' knowledge and skills regarding the clinical problem-solving process in less than two hours. Students were requested to complete a questionnaire throughout this time period in order to obtain data on demographic variables.

Part 2: The delivery of the intervention

After one week, the training instruction began and lasted four weeks (a total of 10 training hours). Throughout the week, tutorials were offered and included problem-based learning strategies as well as appropriate hands-on and small group activities.

This section of the study introduced students to the clinical problem-solving process. This section is divided into three phases, as follows:



Phase I

Through didactic lectures, this phase was aimed to familiarize students with the clinical problem-solving process. This phase involved delivering a series of didactic lectures via Microsoft Teams® on pharmaceutical and disease information resources. The following websites were discussed and used throughout the course: accesspharmacy.mhmedical.com, <https://online.epocrates.com>, www.Drugs.com, www.elm.jo and www.medscape.com. To get the essential information, participants must consult at least two of these sources. Additionally, during this time period, pharmacy students were taught to decision-making and professional communication abilities.

Phase II

Three skills-based workshops were performed, also via Microsoft Teams®, to facilitate the translation of theory into practice. Workshop participants were exposed to simulated patient case studies and participated in group discussions. The cases involved a variety of various body systems, including cardiovascular, pulmonary, rheumatic, endocrine, and infectious illnesses. Following these cases, the researchers moderated a structured and open discussion. The purpose of these workshops was to help participants transition from theoretical knowledge to application and active decision-making.

The lecturer for this course (a ground researcher) interacted with students via video conference, providing relevant examples, discussions, workshops, and motivational feedback.

Phase III

This part evaluated the competencies acquired by attendance at lectures and workshops on the subject. Students were divided into groups and assigned cases. Each group was assigned a portion of the case study to deliver in a presentation manner.

Part 3: Post-intervention

Following that, cases were discussed and skills in data collection, assessment, care plans and patient education were evaluated over a 5-week period, with each group having two hours of discussion. Then, students' knowledge and skills about the clinical problem-solving process were re-evaluated, as well as their level of satisfaction with the training workshops. Additionally, we compared the final exam results and presentation and case discussion skills of students enrolled in the current interventional tutorial to those who got the standard tutorial the previous semester in the same setting.

Students were evaluated in this study using an oral test, multiple-choice and written exams and tutorial participation. Three lecturers (all with a PhD. in Clinical Pharmacy) and three pharmacists facilitated the tutorial.

An online post-intervention test was administered. Kirkpatrick's Model's first two levels (Reaction and Learning) were utilized to evaluate the new tutorial tool's efficacy. The Kirkpatrick model can be used before, during, and following training to demonstrate the training's worth to an organization or an individual. The model is divided into four levels.⁴⁵ Due to the

cost and difficulty of assessing levels 3 and 4 (Behavior and Results, respectively), only the first two levels (Reaction and Learning) were tested during this course.

Level 1 (Reaction; "to what extent participants react positively to the learning event") was measured using a satisfaction questionnaire that judged pharmacy students' positive reactions to the tutorial. To ensure the questionnaire's clarity and usability, it was face and content validated by the study's researchers. Each statement was evaluated using a five-point Likert scale (strongly agree, agree, neutral, disagree, and strongly disagree).

Two criteria were used to assess Level 2 (Learning; "the extent to which participants acquire the desired knowledge, abilities, and attitudes as a result of their involvement in the learning event"):

1) Acquired knowledge of the clinical problem-solving procedure by students. This section included an exam designed to assess pharmacy students' knowledge of the clinical problem-solving process. Three researchers face and content validated the exam in order to improve the clarity and comprehensibility of the questions. The questionnaire had 30 multiple-choice questions covering various topics such as data collection, assessment, treatment plan development, and patient education.

2) Acquired abilities of students to apply the clinical problem-solving method. During this phase, students were divided into groups of five to seven and asked to analyze a variety of clinical cases of similar complexity.

Part 4: Focus group discussions

Additionally, focus group discussions were held online via Microsoft Teams®. We used a standardized interview protocol. Students who completed all three phases were invited to engage in an online focus group discussion. This phase of the study aims to collect comments from participants regarding the intervention trial. It provided an opportunity for reflection on the intervention and its impact. The ground researcher transcribed, de-identified and translated this conversation from Arabic to English. The ground researcher coded, conceptually analyzed and highlighted emerging topics from the translated discussion. The themes were validated following a separate analysis of the data by the study team (three researchers, all with Ph.D. in clinical pharmacy).

Data analysis

Data was analysed by the Statistical Package for Social Science (SPSS Inc., Chicago, IL, USA) version 24. Mean \pm standard deviation (SD) was used to express the continuous variables and frequencies (percentages) were used to express categorical variables.

Checking for normality was carried out using Shapiro-Wilk test (with $p > 0.05$ indicating a normally distributed continuous variable) or by visual inspection of a Normal Q-Q Plot. Paired t-test was used to evaluate pre-post changes in the knowledge and skills scores; if the data was not normally distributed, Wilcoxon sign rank test was used instead. Independent t-test



was used to compare the scores between the students enrolled in the current interventional tutorial and other students who received the traditional tutorial in the preceding semester. For all statistical analysis, $p < 0.05$ was considered statistically significant.

RESULTS

Among the 155 students who registered to in the Clinical Pharmacy and Therapeutics Lab module, 129 participated in this study and completed the educational training (response rate 83.2%), with a mean age (\pm SD) of 23.0 (\pm 1.9) years. Female represented 74.4% of the study sample ($n=96$). Regarding year level, the majority ($n=126$, 97.7%) of the participants were in their fifth (final) year of the undergraduate pharmacy program. Whilst, 2.3 % of the participants were fourth year students who had completed other prerequisite modules. None of the participating students was repeating this module due to a previous failure. As stated by the students, around half of the students (53.5%) had attended a course/workshop about the clinical problem-solving process (Table 1).

Parameters	n (%)	Mean (SD)
Gender		
Male	33 (25.6)	
Female	96 (74.4)	
Age (years)		23.00 (\pm 1.90)
Year of Study		
Fourth year	3 (2.3)	
Fifth year	126 (97.7)	
Attended a course/workshop about clinical problem-solving process		
Yes	69 (53.5)	
No	60 (46.5)	

As shown in Table 2, the majority of students (77.5%) preferred to solve the clinical case as a group rather than individually. Most of the students believed that working as a group improves their skills in problem solving with a mean (\pm SD) of 4.07/5 (\pm 1.11). Also, most of students agreed that working as a group improved their skills in discussion and decision making with means (\pm SD) of 4.17 (\pm 0.99) and 4.16/5 (\pm 0.91), respectively. In the context of the obstacles that the students could face when they deal with each other as a group, the highest proportion of the students (28.7 %) thought that the time coordination within the group is the most faced obstacle. However, the second highest proportion of the students (25.6 %) did not find any obstacles when dealing with their group.

The improvement in students' knowledge and skills following the intervention was assessed and presented in Table 3 and the results showed that there was a significant improvement in the students' knowledge and skills.

Parameters	n (%)
Prefer to solve the case individually	29 (22.5)
Prefer to solve the case as a group	100 (77.5)
Working as a group improves your skills in problem-solving	
Strongly agree (5)	54 (41.9)
Agree (4)	51 (39.5)
Neutral (3)	11 (8.5)
Disagree (2)	5 (3.9)
Strongly disagree (1)	8 (6.2)
Mean (SD)	4.07 (1.11)
Median	4.00
Working as a group improves your skills in discussion	
Strongly agree (5)	55 (42.6)
Agree (4)	56 (43.4)
Neutral (3)	9 (7.0)
Disagree (2)	3 (2.3)
Strongly disagree (1)	6 (4.7)
Mean (SD)	4.17 (0.99)
Median	4.00
Working as a group improves your skills in decision making	
Strongly agree (5)	52 (40.3)
Agree (4)	54 (41.9)
Neutral (3)	17 (13.2)
Disagree (2)	3 (2.3)
Strongly disagree (1)	3 (2.3)
Mean (SD)	4.16 (0.91)
Median	4.00
The most obstacle you face when dealing with your group	
Time coordination	37 (28.7)
Communication skills and ways	13 (10.1)
Conflicts	4 (3.1)
Convincing others	2 (1.6)
All the mentioned	22 (17.1)
Others	18 (14.0)
No obstacle was found	33 (25.6)

A paired-samples t-test was used to determine whether there was a statistically significant mean difference between the students' exam scores (case discussion and report) after the tutorial compared to those scores before the tutorial. As assessed by inspection of a boxplot, there were no outliers in the data. The differences between the pre- and post-tutorial scores were normally distributed, as assessed by visual inspection of a Normal Q-Q Plot. The participants achieved higher score following the tutorial (14.77 ± 2.60) than the baseline (9.38 ± 5.71), with a statistically significant difference of 5.39 (95% CI, 4.53 to 6.24), $t(128) = 12.46$, $p < 0.001$ (Table 3).



Score type	Pre-training	Post-training	The mean difference	95% CI of the Difference		t	df	P value
	Mean (SD)			Lower	Upper			
A paired-samples t-test								
Case discussion and report score	9.38 (5.71)	14.77 (2.60)	5.39 (3.27)	4.53	6.24	12.458	128	<0.001
Students' self-score								
Data Collection	53.07 (21.58)	81.00 (13.87)	27.93 (18.63)	24.68	31.18	17.026	128	<0.001
Assessment	48.56 (24.57)	79.75 (13.60)	31.19 (22.08)	27.35	35.04	16.049	128	<0.001
Care Plan	41.412 (25.10)	78.80 (15.23)	37.39 (23.07)	33.37	41.41	18.405	128	<0.001
Patient Education	57.98 (20.80)	82.98 (13.81)	24.99 (18.02)	21.85	28.13	15.750	128	<0.001
Follow-Up	51.18 (25.28)	79.05 (16.35)	27.88 (21.19)	24.19	31.57	14.944	128	<0.001
An independent-samples t-test								
	Traditional tutorial	Interventional tutorial						
Final exam score	25.54 (7.46)	43.05 (4.25)	17.51 (6.01)	-19.36	-15.66	-21.43	203	< 0.001
Case discussion and presentation evaluation score	16.16 (4.11)	16.72 (2.54)	0.56 (3.11)	-1.47	0.36	-1.21	203	0.23

A paired-samples t-test was used to determine whether there was a statistically significant mean difference between the students' self-reported scores of five skills; data collection, assessment, care plan, patient education, follow-up after the tutorial compared to those scores before the tutorial. There were no outliers in the data. The differences between the pre- and post-tutorial scores of the five parts were normally distributed. As shown in Table 3, all the evaluated five skills were improved significantly following the tutorial, with the highest improvement were reported for the care plan, followed by assessment, data collection, follow-up and finally patient education ($p < 0.005$).

An independent-samples t-test was run to determine if there were differences in the final exam score (out of 50) between the students enrolled in the interventional tutorial ($n = 129$) and other students who received the traditional tutorial in the preceding semester ($n = 76$). The scores for each level of the groups were normally distributed, as assessed by Shapiro-Wilk's test ($p > 0.05$), and there was homogeneity of variances, as assessed by Levene's test for equality of variances ($p = 0.160$). The intervention group achieved higher score (43.05 ± 4.25) than the control group (25.54 ± 7.46), a statistically significant difference of 17.51 (95% CI, -19.36 to -15.66), $t(203) = -21.43$, $p < 0.001$ (Table 3).

An independent-samples t-test was run to determine if there were differences in the case discussion and presentation evaluation score (out of 20) between the students enrolled in the interventional tutorial ($n = 129$) and other students who received the traditional tutorial in the preceding semester ($n = 76$). The intervention group achieved higher score (16.72 ± 2.54) than the control group (16.16 ± 4.11), a statistically insignificant difference of 0.56 (95% CI, -1.47 to 0.36), $t(203) = -1.21$, $p = 0.23$ (Table 3).

Table 4 represents the number of treatment related problems

detected by students pre-training (TRP-1), and post-training using two approaches; TRP-2 using the skills learned during the tutorial (indication, effectiveness, safety, patient, miscellaneous) or TRP-3 using the previous validated checklist.⁴⁶ Based on the means comparison, both post-tutorial approaches showed increase in the number of TRPs detected by students with means \pm SD of 5.80 ± 5.11 and 5.53 ± 6.48 , respectively, compared to pre-tutorial 4.51 ± 3.03 . The paired difference in TRP-3 was insignificant compared to TRP-1. However, there was a significant increase in the number of TRPs detected by students in TRP-2 (1.29) (95% CI, 0.55 to 2.03), $t(127) = 3.46$, $p < 0.001$ (Table 4). The easiness and usefulness of TRP-2 and TRP-3 were assessed. 54.3% of the students thought that TRP-2 is easier than TRP-3. Moreover, most of the students (68.2%) thought that TRP-2 is more useful than TRP-3.

The student's satisfaction with the training tutorial about the clinical problem-solving process (post educational tutorial) was assessed (Table 5). Most of the students demonstrated that the educational training tutorial enhanced their learning as overall with a mean \pm SD of $4.22/5 \pm 0.75$. Furthermore, most of the students found that the patient education instructions and the drug information resource instructions were well administrated (mean = 4.27 ± 0.81 and 4.29 ± 0.74 , respectively). Also, the bulk

	TRP_1	TRP_2	TRP_1	TRP_3
Mean	4.51	5.80	4.51	5.53
SD	3.03	5.11	3.03	6.48
Paired differences	1.29 (4.22)		1.01 (6.64)	
95% CI of the differences	0.55 to 2.03		-0.15 to 2.17	
t	$t(127) = 3.46$		$t(127) = 1.72$	
p value	$p = 0.001$		$p = 0.088$	



Table 5. Students Satisfaction with the educational training tutorial about the clinical problem-solving process (post educational tutorial) (N = 129)								
No	Statements	5	4	3	2	1	Mean (SD)	Median (IQR)
Part A:								
1	The educational tutorial stimulated my interest in Clinical problem-solving process.	37 (28.7)	75 (58.1)	13 (10.1)	3 (2.3)	1 (0.8)	4.12 (0.74)	4 (0)
2	Methods of Clinical problem-solving process introduced in this educational tutorial are beneficial.	37 (28.7)	66 (51.2)	22 (17.1)	4 (3.1)	0	4.05 (0.76)	4 (0)
3	I enjoyed participating in this education training.	35 (27.1)	56 (43.4)	29 (22.5)	7 (5.4)	2 (1.6)	3.89 (0.92)	4 (1)
4	The educational tutorial was easy to understand.	28 (21.7)	42 (32.6)	34 (26.4)	23 (17.8)	2 (1.6)	3.55 (1.07)	4 (1)
5	The educational tutorial made me understand the concept of Clinical problem-solving process.	45 (34.9)	60 (46.5)	21 (16.3)	3 (2.3)	0	4.14 (0.77)	4 (0)
6	The educational tutorial helped me understand the importance of Clinical problem-solving process.	45 (34.9)	52 (40.3)	30 (23.3)	2 (1.6)	0	4.09 (0.80)	4 (0)
7	The educational tutorial enhanced my learning.	50 (38.8)	60 (46.5)	16 (12.4)	3 (2.3)	0	4.22 (0.75)	4 (0)
8	Wide knowledge area covered.	47 (36.4)	49 (38.0)	29 (22.5)	4 (3.1)	0	4.08 (0.84)	4 (0)
9	Wide range of clinical skills covered.	53 (41.1)	48 (37.2)	25 (19.4)	3 (2.3)	0	4.17 (0.82)	4 (0)
10	The educational tutorial well structured & sequenced.	36 (27.9)	57 (44.2)	30 (23.3)	4 (3.1)	2 (1.6)	3.94 (0.88)	4 (0)
11	Student aware of level of data and skills needed.	46 (35.7)	51 (39.5)	27 (20.9)	4 (3.1)	1 (0.8)	4.06 (0.87)	4 (0)
Part B:								
1	DI resources instructions well administered.	55 (42.6)	59 (45.7)	13 (10.1)	1 (0.8)	1 (0.8)	4.29 (0.74)	4 (0)
2	DI resources skills are beneficial.	54 (41.9)	54 (41.9)	9 (7.0)	3 (2.3)	0	4.30 (0.70)	4 (1)
3	Data Collection part instructions well administrated.	41 (31.8)	56 (43.4)	28 (21.7)	4 (3.1)	0	4.04 (0.81)	4 (0)
4	Data Collection part is clear.	42 (32.6)	48 (37.2)	30 (23.3)	8 (6.2)	1 (0.8)	3.95 (0.94)	4 (1)
5	Data Collection part is beneficial.	53 (41.1)	57 (44.2)	15 (11.6)	4 (3.1)	0	4.23 (0.78)	4 (0)
6	Medication's assessment template instructions well administrated.	43 (33.3)	54 (41.9)	25 (19.4)	5 (3.9)	2 (1.6)	4.02 (0.91)	4 (0)
7	Medication's assessment part is clear.	42 (32.6)	51 (39.5)	24 (18.6)	10 (7.8)	2 (1.6)	3.94 (0.98)	4 (0)
8	Medication's assessment part is beneficial.	58 (45.0)	44 (34.1)	22 (17.1)	3 (2.3)	2 (1.6)	4.19 (0.91)	4 (0)
9	The educational training helped me to be able to identify treatment-related problems (TRPs).	54 (41.9)	49 (38.0)	20 (15.5)	3 (2.3)	3 (2.3)	4.15 (0.93)	4 (0)
10	Care plan template instructions well administrated.	43 (33.3)	54 (41.9)	26 (20.2)	3 (2.3)	3 (2.3)	4.02 (0.92)	4 (0)
11	Care plan part is clear.	46 (35.7)	50 (38.8)	22 (17.1)	9 (7.0)	2 (1.6)	4.00 (0.98)	4 (0)
12	Care plan part is beneficial.	57 (44.2)	53 (41.1)	14 (10.9)	5 (3.9)	0	4.26 (0.80)	4 (0)
13	Patient education instructions well administrated.	60 (46.5)	48 (37.2)	17 (13.2)	4 (3.1)	0	4.27 (0.81)	4 (1)
14	Patient education part is clear.	58 (45.0)	48 (37.2)	20 (15.5)	3 (2.3)	0	4.25 (0.80)	4 (1)
15	Patient education part is beneficial.	53 (41.1)	58 (45.0)	15 (11.6)	3 (2.3)	0	4.25 (0.75)	4 (0)
16	It is beneficial to include the skills learned in different courses like therapeutics, OTC, training, clinical biochemistry, DI etc.	56 (43.4)	43 (33.3)	20 (15.5)	8 (6.2)	2 (1.6)	4.11 (0.99)	4 (0)
Part C:								
1	Hospital case evaluation was fair.	18 (14.0)	60 (46.5)	32 (24.8)	18 (14.0)	1 (0.8)	3.59 (0.92)	4 (1)

2	Final exam was fair.	47 (36.4)	53 (41.1)	25 (19.4)	4 (3.1)	0	4.11 (0.82)	4 (0)
3	Exams well administered.	38 (29.5)	64 (49.6)	25 (19.4)	2 (1.6)	0	4.07 (0.74)	4 (0)
4	Exams very stressful.	32 (24.8)	34 (26.4)	32 (24.8)	18 (14.0)	13 (10.1)	3.42 (1.28)	4 (1)
5	Exams well structured & sequenced.	38 (29.5)	52 (40.3)	33 (25.6)	5 (3.9)	1 (0.8)	3.94 (0.88)	4 (0)
6	Personality, ethnicity, and gender do not affect scores.	63 (48.8)	36 (27.9)	23 (17.8)	5 (3.9)	2 (1.6)	4.19 (0.97)	4 (0)

5: strongly agree; 4: agree; 3: neutral; 2: disagree; 1: strongly disagree.

of students found that the patient education part was clear and beneficial with means \pm SDs of 4.25 ± 0.80 and 4.25 ± 0.80 , respectively. Moreover, the majority of students found that the care plan part was beneficial (mean= 4.26 ± 0.80), the data collection part is beneficial as well (mean= 4.23 ± 0.78), and the drug information resources skills were beneficial (mean= 4.30 ± 0.70). Regarding the exam, most of students thought that it was fair (mean= 4.11 ± 0.82), in addition, that the personality, ethnicity and gender did not affect the scores of the exam (mean= 4.19 ± 0.97).

Finally, the majority of students were satisfied with the overall training process and stated a high evaluation score out of 10 (mean = 7.93 ± 1.42 , median = 8.00) (Figure 1).

In addition to the positive changes observed in the post-surveys, focus-group discussions and students' opinions appeared to endorse findings of enhanced understanding of clinical problem-solving skills. Discussions were transcribed, translated and analysed by the ground researcher and separately analysed by the research team.

Students identified the need for and the impact of such courses on pharmacy practice in Jordan. They repeatedly described the course as "interesting" and "thought-provoking", expressing

the novelty of this topic to them, and depicted this teaching approach as "beneficial". Participants in this study commented positively about workshop discussions utilized, explaining that this approach facilitates the transforming of the theoretical part of the intervention to a practical interactive model, allowing students to learn from each other.

Students who participated in this study considered this training methods useful and demonstrated enhanced confidence in handling clinical cases that might be encountered in practice.

Most participants suggested incorporating such a course in an earlier year of study, then slightly increasing the level of intensity and complexity of information each time repeated across each year of the five years of the BPharm curriculum. Therefore, learning materials are disseminated across all the degree stages in order for pharmacists to practice in a less stressful environment with more confidence.

DISCUSSION

Nowadays, the pharmacist's role has developed dramatically, and it is anticipated that pharmacists will play an important role in providing PC services. Previously, pharmacists have not been




Figure 1. The Overall students' evaluation for the educational tutorial tool (10: the best, 0: the worst). Mean = 7.93 ± 1.42 , median = 8.00.



directly concerned with gathering a good medication history in most hospitals.⁴⁷ This may be due to the insufficient information of the process, the lack of awareness, imprecise understanding of who has the obligation for delivering PC services, and shifting responsibilities to other healthcare providers for administering the service.^{48,49} The lack of knowledge, skills and awareness about the service was among the barriers for implementing PC services. Therefore, this study was designed to solve this gap in learning the clinical problem-solving process at pharmacy school years by the development of an innovative learning tool.

Few studies have been conducted to determine the effect of educational programs on students' ability to obtain medication histories and to identify and resolve TRPs.⁵⁰⁻⁵² These studies have focused on incorporating an interactive learning exercise using simulation^{50,51} and the use of a short duration video tutorial⁵² to educate students about medication reconciliation and the results of these studies indicated that students' knowledge and skills about medication reconciliation and identifying discrepancies were improved following the educational intervention.⁵⁰⁻⁵² However, medication reconciliation and finding discrepancies are only one part of PC principle. Accordingly, the aim of this study was to develop, implement and evaluate the effectiveness of another teaching method, an online tutorial, to educate pharmacy students about the clinical problem-solving process and how to retrieve patient data from medical files and developing a patient specific pharmaceutical care plan for the identification, resolution and prevention of TRPs that enable them to practice PC and improve decision making process.

Pharmacy education provided at the level of pharmacy schools is considered as one of the most crucial elements in the development of professional pharmacists.^{50,53,54} Pharmacy educators have the responsibility to provide students with adequate and comprehensive training and educational tools to ensure their ability to provide the PC service to their patients.^{50,53,54} Therefore, it is important to supply students with the proper skills in a way that the pharmacy educators guarantee that they obtained appropriate skills during their training and to be able to provide PC correctly.^{50,53,54}

New techniques are getting attention in the field of education, involving the usage of online communication and videos.^{55,56} Visual teaching tools have been shown to improve students' comprehension and learning.⁵⁶⁻⁵⁸ However, their effectiveness in pharmacy education is not fully known.^{56,59,60} Problem-based learning still remains the most innovative medical education method as it is characterized by self-directed learning using simulated real-life scenarios as the learning programme in addition of being student-centered learning.⁶¹

The findings of this interventional study demonstrate that this training method significantly improved the total knowledge and skills scores about clinical problem-solving process. The participants achieved higher examiner scores (case discussion and report) following the tutorial compared to those scores before the tutorial, with a statistically significant difference, $p < 0.001$. Also, by investigating the students' self-reported scores, all the evaluated five skills were improved significantly following the tutorial, with the highest improvement were reported for

the care plan, followed by assessment, data collection, follow-up, and finally patient education ($p < 0.005$). We compared the final exam score between the students enrolled in the interventional tutorial and other students who received the traditional tutorial in the preceding semester. The intervention group achieved higher score than the control group, $p < 0.001$. Unlike a previous recent study which evaluated the influence of the training module immediately after conducting the workshop,⁵² we evaluated the effectiveness after several weeks which ensure measuring the long-term effect of this educational tutorial.

The study's findings were consistent with those of prior research done in the United Kingdom and New Zealand in certain aspects.^{56,62} The researchers in these studies discovered that visual techniques of instruction produced positive effects.⁵⁶ For instance, videos demonstrating several practical skills and visual aids represented in Magnetic Resonance Imaging (MRI) images were effective and improved the understanding of students about certain pharmaceutical concepts.⁶² Moreover, enhancement of innovative and interactive learning experience of students were observed.^{56,62}

In terms of students' ability to detect TRPs, students were able to identify more TRPs following the instruction. As a result, it may be assumed that students who participated in this study were able to put their newly acquired information and abilities to use.

When students' perceived satisfaction with their training strategy was evaluated, the majority expressed pleasure with the whole training process and stated that the educational tool helped them acquire the notion of clinical problem-solving. This results was consistent with that of Zhang et al., who demonstrated that using videos in the learning process resulted in a higher level of student satisfaction than other techniques.⁶³ Thus, it can be advised and encouraged that visual educational videos be incorporated into the learning process. Additionally, the utilization of online tutorials as a teaching approach benefited from the fact that they were available at any time. In contrast to typical lectures, students can listen to the information multiple times and refer to the recordings if they missed anything.

Upon using sound methods of educational delivery, educational intervention trials have demonstrated an enhancement of the pharmacy practice skills.^{64,65} Moreover, implementing such pedagogically-framed training has shown an improvement in the quality of patient-care in addition to enhancing the clinical problem-solving process. Incorporating role-play, team-based learning, and stimulated-patient encounters as teaching methods have been highlighted in the literature and proved to affect participants' learning experience positively.^{31,64,66-69}

Literature has identified students' aspiration for instructors to adapt to students' needs by implementing meaningful examples to invoke discussions related to practice and in doing so motivate students to do their best.⁷⁰ The instructor of this course immersed students in the online course via video conferencing, providing related examples, discussions, workshops and motivational feedback. Students who



participated in this study considered these methods useful and demonstrated enhanced confidence in handling clinical cases that might be encountered in practice. Hence these approaches of teaching were selected to be incorporated as components supporting the traditional didactic lectures component at baseline, in our course, with good outcomes for the students.

For the first time, this study explored the implementation and evaluation of customised educational intervention that aims to enhance students' knowledge and skills in clinical problem-solving process in Jordan. The findings have demonstrated the remarkable positive impact of this educational intervention on decision-making skills of participating pharmacy students. The study highlighted in particular, a significant increase in participants' confidence level in decision-making after joining the course. The students' feedback, also, instruct us about the perceived need/utility, impact and suggestions to be considered in the future regarding the delivery of this course.

Furthermore, this study demonstrated that the delivered standard education in the clinical pharmacy and therapeutics tutorials was quite sufficient to provide the pharmacy students with the needed knowledge and skills to demonstrate the clinical problem-solving process. Such interventions were considered effective in helping students attain basic skills such as teamwork, peer assessment, communication and critical evaluation.

To serve the patients requirements, ample opportunity should be given to the students to develop their clinical knowledge foundation and improve their communication skills effectively. It is equally important to install in students' positive attitudes and motivation to provide PC.³⁷ This study highlighted the need for incorporating such courses in the undergraduate curricula in all universities in Jordan.

The findings of this study need to be interpreted with caution because of some limitations. First of all, the effectiveness on real practice was not assessed, and students' skills in the clinical problem-solving process were evaluated only using few cases scenario that applies to the hospital/clinic settings. To resolve many of the obstacles recognized in this study, facing pharmacy students with real cases and providing them with a practical educational approach to deliver counselling to these real patients are required.^{71,72} Further investigation is still needed to find ways, with enhanced educational and training strategies, that overcome the identified barriers revealed in this study. These ways should be able to improve students' confidence and competency in patient counselling once they start practicing pharmacy.

Also, our study was conducted only among one cohort of students from one university which may limit the generalizability of the results. However, no major differences can be found in the other schools of pharmacy that can limit the generalization of the findings of this study and much can be learned from this initial trial.

This study was conducted in unprecedented COVID-19 pandemic circumstances, which mandated lockdowns and travel restrictions, thereby obligating the construct of the

educational intervention to be in an online format and adding some complexity to delivery and evaluation. Online teaching has often been improvised rapidly in such circumstances.⁷³ Yet despite the challenges, the delivery of this course conformed with aspects identified as essential components for efficient information delivery modes, yielding positive student outcomes focusing on competency development.⁷³

Final limitation of this study is that students might have responded in a socially undesirable manner in the post-intervention survey, which we addressed by thoroughly inspecting responses and discarding irregular or "bogus data".

CONCLUSION

This study explored the implementation and evaluation of an educational medical informatics tutorial that aims to enhance students' knowledge and skills in clinical problem-solving process in Jordan. The results of this interventional study indicate that this educational intervention achieved a significant positive impact of on decision-making skills of participating pharmacy students. The study highlighted in particular, a significant increase in participants' confident level in decision-making after joining the course. Furthermore, such intervention was considered effective in helping students attain basic skills such as teamwork, peer assessment, communication and critical evaluation.

Effective medical care improvement and PC requires the collaboration of healthcare providers in ensuring the accurate use of medications through transitions in care. Among other healthcare professionals, pharmacists as medication experts have a vital role contributing in the success of process implementation. Pharmacy educators are liable for preparing pharmacy student to be ready to implement PC in their future practice.

ACKNOWLEDGEMENTS

We would like to thank every participant and volunteer who participated in this study and Dalal Alnatour, PharmD, Dima Al-awajan, PharmD, Wafa Alhourani, BS.

FUNDING

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

CONFLICTS OF INTEREST

All Authors declare no conflict of interest related to this article.

ABBREVIATIONS

DI: drug information
OTC: over the counter
PC: pharmaceutical care
TRPs: treatment-related problems



References

1. Bradley C. The future role of pharmacists in primary care: *British Journal of General Practice*. *Br J Gen Pract*. 2009;59(569):891-2. <https://doi.org/10.3399/bjgp09X473105>
2. Toklu HZ, Hussain AJJoYP. The changing face of pharmacy practice and the need for a new model of pharmacy education. *J Young Pharm*. 2013;5(2):38-40. <https://doi.org/10.1016/j.jyp.2012.09.001>
3. Routhead E, Semple SJ, Vitry AJJoPP. Pharmaceutical care services: a systematic review of published studies, 1990 to 2003, examining effectiveness in improving patient outcomes. 2005;13:53-70.
4. Hepler CDJPTJoHP, Therapy D. Clinical pharmacy, pharmaceutical care, and the quality of drug therapy. *Pharmacotherapy*. 2004;24(11):1491-1498. <https://doi.org/10.1592/phco.24.16.1491.50950>
5. Aburuz S. The case for evidence-based pharmaceutical care. *Res. Social Adm. Pharm*. 2015;11(3):e146-147. <https://doi.org/10.1016/j.sapharm.2009.06.001>
6. Alsayed A, El Hajji FD, Al-Najjar MA, et al. Patterns of antibiotic use, knowledge, and perceptions among different population categories: A comprehensive study based in Arabic countries. *Saudi Pharmaceutical Journal*. 2022.
7. Khader H, Hasoun LZ, Alsayed A, et al. Potentially inappropriate medications use and its associated factors among geriatric patients: a cross-sectional study based on 2019 Beers Criteria. *Pharmacia*. 2021;68:789-795.
8. Kruijtbosch M, Göttgens-Jansen W, Floor-Schreuderling A, et al. Moral dilemmas of community pharmacists: a narrative study. *Int J Clin Pharm*. 2018;40(1):74-83. <https://doi.org/10.1007/s11096-017-0561-0>
9. Alsayed AR, Hasoun L, Alnatour D, et al. Perspectives of the community in the developing countries toward telemedicine and pharmaceutical care during the COVID-19 pandemic. *Pharm Pract*. 2022;20(1):2618. <https://doi.org/10.18549/PharmPract.2022.1.2618>
10. Hepler CD, Strand LM. Opportunities and responsibilities in pharmaceutical care. *Am J Hosp Pharm*. 1990;47(3):533-543.
11. Allemann SS, van Mil JF, Botermann L, et al. Pharmaceutical care: the PCNE definition 2013. *Int J Clin Pharm*. 2014;36(3):544-555. <https://doi.org/10.1007/s11096-014-9933-x>
12. Shao H, Chen G, Zhu C, et al. Effect of pharmaceutical care on clinical outcomes of outpatients with type 2 diabetes mellitus. *Patient Prefer Adherence*. 2017;11:897. <https://doi.org/10.2147/PPA.S92533>
13. Parajuli DR, Franzon J, McKinnon RA, et al. Role of the pharmacist for improving self-care and outcomes in heart failure. *Curr Heart Fail Rep*. 2017;14(2):78-86. <https://doi.org/10.1007/s11897-017-0323-2>
14. Shanmugam S, Varughese J, Nair MAS, et al. Pharmaceutical care for asthma patients: A Developing Country's Experience. *J Resp Pharm Pract*. 2012;1(2):66. <https://doi.org/10.4103/2279-042X.108373>
15. Cazarim MdS, de Freitas O, Penaforte TR, et al. Impact assessment of pharmaceutical care in the management of hypertension and coronary risk factors after discharge. *PloS One*. 2016;11(6):e0155204. <https://doi.org/10.1371/journal.pone.0155204>
16. Liekweg A, Westfeld M, Braun M, et al. Pharmaceutical care for patients with breast and ovarian cancer. *Support Care Cancer*. 2012;20(11):2669-2677. <https://doi.org/10.1007/s00520-012-1385-z>
17. Kousar R, Murtaza G, Azhar S, et al. Randomized controlled trials covering pharmaceutical care and medicines management: a systematic review of literature. *Res Social Adm Pharm*. 2018;14(6):521-539. <https://doi.org/10.1016/j.sapharm.2017.06.008>
18. Poudel RS, Piryani RM, Shrestha S, et al. Benefit of hospital pharmacy intervention on the current status of dry powder inhaler technique in patients with asthma and COPD: a study from the Central Development Region, Nepal. *Integr Pharm Res Pract*. 2017;6:7-13. <https://doi.org/10.2147/IPRP.S119202>
19. Berenguer B, La Casa C, de la Matta MJ, et al. Pharmaceutical care: past, present and future. *Curr Pharm Des*. 2004;10(31):3931-3946. <https://doi.org/10.2174/1381612043382521>
20. Obreli-Neto PR, Guidoni CM, de Oliveira Baldoni A, et al. Effect of a 36-month pharmaceutical care program on pharmacotherapy adherence in elderly diabetic and hypertensive patients. *Int J Clin Pharm*. 2011;33(4):642-649. <https://doi.org/10.1007/s11096-011-9518-x>
21. Clifford RM, Davis WA, Batty KT, et al. Effect of a pharmaceutical care program on vascular risk factors in type 2 diabetes: the Fremantle Diabetes Study. *Diabetes Care*. 2005;28(4):771-776. <https://doi.org/10.2337/diacare.28.4.771>
22. Bunting BA, Cranor CWJJoTAPA. The Asheville Project: long-term clinical, humanistic, and economic outcomes of a community-based medication therapy management program for asthma. *J Am Pharm Assoc*. 2006;46(2):133-147. <https://doi.org/10.1331/154434506776180658>
23. Cranor CW, Bunting BA, Christensen DB. The Asheville Project: long-term clinical and economic outcomes of a community pharmacy diabetes care program. *J Am Pharm Assoc*. 2003;43(2):173-184. <https://doi.org/10.1331/108658003321480713>
24. May JRJAJoHP. Barriers to pharmaceutical care in the acute care setting. 1993;50:1608-1611.
25. Roberts AS, Benrimoj SI, Chen TF, et al. Practice change in community pharmacy: quantification of facilitators. *Ann Pharmacother*. 2008;42(6):861-868. <https://doi.org/10.1345/aph.1K617>
26. McDonough RP, Rovers JP, Currie JD, et al. Obstacles to the implementation of pharmaceutical care in the community setting. 1998;38:87-95.
27. Ngorsuraches S, Li SCJAjoh-sp. Thai pharmacists' understanding, attitudes, and perceived barriers related to providing pharmaceutical care. *Am J Health Syst Pharm*. 2006;63(21):2144-2150. <https://doi.org/10.2146/ajhp060054>



28. Uema SA, Vega EM, Armando PD, et al. Barriers to pharmaceutical care in Argentina. *Pharm World Sci.* 2008;30(3):211-215. <https://doi.org/10.1007/s11096-007-9167-2>
29. Fang Y, Yang S, Feng B, et al. Pharmacists' perception of pharmaceutical care in community pharmacy: a questionnaire survey in Northwest China. *Health Soc Care Community.* 2011;19(2):189-197. <https://doi.org/10.1111/j.1365-2524.2010.00959.x>
30. Education TACfP. Accreditation Standards and Key Elements for the Professional Program in Pharmacy Leading to the Doctor of Pharmacy Degree. Standards 2016. Vol 2020: Accreditation Council for Pharmacy Education. 2016.
31. Medina MS, Plaza CM, Stowe CD, et al. Center for the Advancement of Pharmacy Education 2013 educational outcomes. *Am. J. Pharm. Educ.* 2013;77(8):162-162. <https://doi.org/10.5688/ajpe778162>
32. Accreditation Standards and Key Elements for the Professional Program in Pharmacy Leading to the Doctor of Pharmacy Degree. Standards 2016. Vol 2019: Accreditation Council for Pharmacy Education. 2016.
33. Hatwig C, Crane V, Hayman JJTihpm. A pharmaceutical care challenge: recruiting, training, and retaining pharmaceutical care practitioners. *Top Hosp Pharm Manage.* 1993;13:1-10.
34. Kane MPJAJoPE. Solving Drug-Related Problems in the Professional Experience Program. 1993;57:347-351.
35. Briceland LL, Hamilton RAJAJHP. Skk. 1993;50:294-296.
36. Chisholm MA, Hawkins DWJJoPT. Analysis of pharmacotherapy recommendations provided by doctor of pharmacy clerkship students. 1996;5:3-12.
37. Chisholm MA, Wade WEJAjoh-sp. Factors influencing students' attitudes toward pharmaceutical care. *Am J Health Syst pharm.* 1999;56:2330-2335. <https://doi.org/10.1093/ajhp/56.22.2330>
38. Chisholm MA, Wade WEJAJoPE. Using actual patients in the classroom to develop positive student attitudes toward pharmaceutical care. 1999;63:296-299.
39. Kennedy DT, Ruffin DM, Goode JVR, et al. The Role of Academia in Community-Based Pharmaceutical Care. *Pharmacotherapy.* 1997;17:1352-1356.
40. Schommer JC, Cable GLJAJoPE. Current status of pharmaceutical care practice: strategies for education. 1996;60:36-41.
41. Hammour KA, Farha RA, Basheti I. Hospital pharmacy medication reconciliation practice in Jordan: perceptions and barriers. *J Eval Clin Pract.* 2016;22(6):932-937. <https://doi.org/10.1111/jep.12565>
42. Ramjaun A, Sudarshan M, Patakfalvi L, et al. Educating medical trainees on medication reconciliation: a systematic review. *BMC Med. Educ.* 2015;15:33. <https://doi.org/10.1186/s12909-015-0306-5>
43. Boockvar KS, Carlson LaCorte H, Giambanco V, et al. Medication reconciliation for reducing drug-discrepancy adverse events. *Am J Geriatr Pharmacother.* 2006;4(3):236-243. <https://doi.org/10.1016/j.amjopharm.2006.09.003>
44. Sluisveld NV, Zegers M, Natsch S, et al. Medication reconciliation at hospital admission and discharge: insufficient knowledge, unclear task reallocation and lack of collaboration as major barriers to medication safety. *BMC Health Serv Res.* 2012;12:170. <https://doi.org/10.1186/1472-6963-12-170>
45. Smidt A, Balandin S, Sigafos J, et al. The Kirkpatrick model: A useful tool for evaluating training outcomes. *Journal of Intellectual & Developmental Disability.* 2009;34(3):266-274. <https://doi.org/10.1080/13668250903093125>
46. AbuRuz SM, Bulatova NR, Yousef AMJPW, et al. Validation of a comprehensive classification tool for treatment-related problems. *Pharm world Sci.* 2006;28(4):222-232. <https://doi.org/10.1007/s11096-006-9048-0>
47. Bond C, Raehl CL, Franke TJPTJoHP, et al. Clinical pharmacy services, hospital pharmacy staffing, and medication errors in United States hospitals. *Pharmacotherapy.* 2002;22(2):134-147. <https://doi.org/10.1592/phco.22.3.134.33551>
48. Boockvar KS, LaCorte HC, Giambanco V, et al. Medication reconciliation for reducing drug-discrepancy adverse events. *Am J Geriatr Pharmacother.* 2006;4(3):236-243. <https://doi.org/10.1016/j.amjopharm.2006.09.003>
49. Sluisveld NV, Zegers M, Natsch S, et al. Medication reconciliation at hospital admission and discharge: insufficient knowledge, unclear task reallocation and lack of collaboration as major barriers to medication safety. *BMC Health Serv Res.* 2012;12:1-12. <https://doi.org/10.1186/1472-6963-12-170>
50. Lubowski TJ, Cronin LM, Pavelka RW, et al. Effectiveness of a medication reconciliation project conducted by PharmD students. *Am J Pharm Educ.* 2007;71(5):94. <https://doi.org/10.5688/aj710594>
51. Lindquist LA, Gleason KM, McDaniel MR, et al. Teaching medication reconciliation through simulation: a patient safety initiative for second year medical students. *J Gen Intern Med.* 2008;23(7):998-1001. <https://doi.org/10.1007/s11606-008-0567-3>
52. Abu Farha RK, Rashad M, Hasen E, et al. Evaluation of the effect of video tutorial training on improving pharmacy students' knowledge and skills about medication reconciliation. *Pharm Pract.* 2020;18(1):1711. <https://doi.org/10.18549/PharmPract.2020.1.1711>
53. Walker PC, Tucker Jones JN, Mason NA. An advanced pharmacy practice experience in transitional care. *Am J Pharm Educ.* 2010;74(2):20. <https://doi.org/10.5688/aj740220>
54. Lancaster JW, Grgurich PE. Impact of students pharmacists on the medication reconciliation process in high-risk hospitalized general medicine patients. *Am J Pharm Educ.* 2014;78(2):34. <https://doi.org/10.5688/ajpe78234>
55. Fox LM, Pham KH, Dollar M. Using interactive digital images of products to teach pharmaceuticals. *Am J Pharm Educ.* 2007;71(3):58. <https://doi.org/10.5688/aj710358>
56. Curley LE, Wu Z, Svirskis D. Using Technology in Pharmacy Education: Pharmacy Student Performance and Perspectives When Visual Aids Are Integrated Into Learning. *Front Pharmacol.* 2018;9:1062. <https://doi.org/10.3389/fphar.2018.01062>



57. Woolfitt ZJLT, Learning, Sciences TIUoA. The effective use of video in higher education. 2015;1:1-49.
58. Brame CJJCLSE. Effective educational videos: Principles and guidelines for maximizing student learning from video content. *CBE Life Sci Educ*. 2016;15(4):es6. <https://doi.org/10.1187/cbe.16-03-0125>
59. Wilson EA, Park DC, Curtis LM, et al. Media and memory: the efficacy of video and print materials for promoting patient education about asthma. *Patient Educ Couns*. 2010;80(3):393-398. <https://doi.org/10.1016/j.pec.2010.07.011>
60. Von Schantz S, Katajavuori N, Juppo AMJP. The use of video instructions in patient education promoting correct technique for dry powder inhalers: An Investigation on Inhaler-Naïve Individuals. *Pharmacy*. 2018;6(4):106. <https://doi.org/10.3390/pharmacy6040106>
61. Kwan CY. A thorny path: the developmental course of problem-based learning for health sciences education in Asia. *Adv Health Sci Educ Theory Pract*. 2019;24(5):893-901. <https://doi.org/10.1007/s10459-019-09920-6>
62. Hitch G, Williams J, Herbland A, et al. The use of videos in blended learning to enhance students' learning in systems-based patient assessment with development of associated clinical skills: An Analysis. 2014;14.
63. Lindner K, Blosser G, Cunigan KJMPS. Visual versus auditory learning and memory recall performance on short-term versus long-term tests. 2009;15:6.
64. Fejzic J, Barker M. Implementing simulated learning modules to improve students' pharmacy practice skills and professionalism. *Pharm. Pract*. 2015;13(3):583. <https://doi.org/10.18549/PharmPract.2015.03.583>
65. DeMella C, Donohoe K, Morgan L, et al. Implementation of an online ethics course into a pharmacy curriculum. *Currents in Pharmacy Teaching and Learning*. 2017;9(5):835-840. <https://doi.org/10.1016/j.cptl.2017.05.001>
66. Al-Aqeel SA. Pharmacy Students Feedback on the use of Role-play in Teaching Ethics. *Pharmacy Education*. 2013;13.
67. Sutton B, Bearman M, Jolly B, et al. Simulated Learning Environments Medical Curriculum Report. Health Workforce Australia. 2010.
68. Lundeberg MA, Yadav A. Assessment of case study teaching: Where do we go from here? Part II. *Journal of College Science Teaching*. 2006;35:8.
69. Hasan S. Teaching ethics to pharmacy students using a team-based learning approach. *Pharmacy Education*. 2015;11.
70. Pinchevsky-Font T, Dunbar S. Best practices for online teaching and learning in health care related programs. *Internet Journal of Allied Health Sciences and Practice*. 2015;13:8.
71. Basheti I. Effect of simulation in training pharmacy students on correct inhaler technique. *Proceedings of International Academic Conferences: International Institute of Social and Economic Sciences*; 2014.
72. Odenina F, Segal R, Helper CJoS, et al. Providing pharmaceutical care in community practice: differences between providers and non-providers of pharmaceutical care. 1995;12:170-180.
73. Rapanta C, Botturi L, Goodyear P, et al. Online University Teaching During and After the Covid-19 Crisis: Refocusing Teacher Presence and Learning Activity. *Postdigital Science and Education*. 2020;2:923-945.

APPENDIX

Table A. Data Collection

1.	Demographic Data:
	<ul style="list-style-type: none"> • Who is the patient? • Age: • Gender: • Education and Occupation:
2.	Current Issues
	<ul style="list-style-type: none"> ❖ Chief Complain: ❖ History of Present Illness: • Location: • Characteristic of symptoms: • Timing (onset, duration, frequency): • Severity: • Factors: Aggravating factors: Alleviating factors: • Environment: • Other symptoms: • Differential diagnosis: ❖ Review of Systems: ❖ Physical Examination:



3. Medical problems
<ul style="list-style-type: none"> • Past, acute and chronic diseases: • Stage / type / class: • Current status: • Duration:
4. Patient History
<ul style="list-style-type: none"> • Surgeries: • Hospitalization: • Vaccination: • Allergies: • Family history: • Social history:
5. Treatments
<ul style="list-style-type: none"> • Drug Details: • Duration: • Time of administration: • Adherence:
6. Tests
<ul style="list-style-type: none"> • Vital signs: • Lab tests: • Other tests:
7. Special Situation
<ul style="list-style-type: none"> • Pediatric, geriatric, pregnancy, breastfeeding, ethnicity. • Limitations (NPO, bedridden, inability to swallow) • Renal impairment • Hepatic impairment • Allergies • Sensitive issues / barrier to communication • Abuse

Table B. Basic calculations

Date	
Weight (kg)	
Height (cm) https://www.feettometres.com/	
BMI [Weight (kg) / Height (m) ²] https://www.nhlbi.nih.gov/health/educational/lose_wt/BMI/bmi-m.htm	
BMI Categories: Underweight = <18.5 Normal weight = 18.5–24.9 Overweight = 25–29.9 Obesity = BMI of 30 or greater	
IBW https://www.mdcalc.com/ideal-body-weight-adjusted-body-weight This formula is only an approximation, and is generally only applicable for people ≥152.4 cm tall. For patients under 152.4 cm, one commonly-used modification is to subtract 1 kg for each 2.54 cm below 152.4 cm.	
Renal impairment (eGFR) https://www.kidney.org/professionals/kdoqi/gfr_calculator eGFR: 90–120 mL/minute/1.73 m ²	

Table C. Assessment of medications

Date	
Drug name (scientific and trade name) / strength / route / frequency	
Medical problem or health care need	



Goals of treatment	
Category 1. PTA, current, or discharge; 2. P, OTC, or CAM 3. PRN; Stat; or regular	
Start date	
Stop date	
Time	
Indication	
Effectiveness	
Safety	
Dosing regimen	
Product preparation (procedures, methods), compatibility and stability	
Drug interactions (drugs, foods, tests)	
Patient (knowledge, adherence)	
Reference	

Abbreviations: PTA: prior to admission; P: Prescribed; OTC: over-the counter; CAM: complementary and alternative medicine; PRN: per registered nurse (as needed).

Table D. Patient Care Plan

Date	
Medical problem or healthcare need	
Goal(s) of Treatment	
Treatment related problem (TRP) [Indication, Effectiveness, Safety, Patient, Others]	
Intervention(s) / Recommendation(s)	



Follow-up (Monitoring and Evaluation) Monitoring parameter(s) (therapeutic or toxic) / Endpoints / Frequency Follow-up (timeframe)	
References	

