

Original Research

Knowledge, practice and attitudes regarding antibiotics use among Lebanese dentists

Hicham MANSOUR, Mireille FEGHALI, Nadine SALEH, Mona ZEITOUNY.

Received (first version): 23-May-2018

Accepted: 20-Aug-2018

Published online: 29-Sep-2018

Abstract

Objectives: Explore antibiotic use, assess conformity with evidence-practice guidelines, and describe knowledge and attitudinal factors among Lebanese dentists.

Methods: National cross-sectional telephonic survey, using a standardized questionnaire addressing demographic, educational and professional data, usual antibiotics prophylactic and curative prescription pattern and influential factors, knowledge concerning antibiotics use in selected patient-populations, and attitude regarding antimicrobial resistance. Analyses used descriptive statistics, and bivariate analysis to observe predictors of higher knowledge.

Results: the overall response rate for the study was around 21%. 322 dentists participated. On average, 17.51% of consultations resulted in antibiotic use; previous antibiotic experience mostly influenced prescriptions (81.3%). Referral of pregnant and lactating women and cardiac patients, when antibiotics are needed, was high (26.9%, 28.5% and 79.4%, respectively). Macrolides were the dominant first-line antibiotics in penicillin allergy (47.4%). Penicillins were most common for pregnant and lactating women. Penicillins (95.0%), 2g (63.9%), and 1 hour pre-procedure (34%) were the main components of prophylaxis for cardiac patients. Prophylactic and curative use varied widely; few dentists exhibited guideline-conform prescriptions. Mean knowledge scores of prophylaxis for cardiac and non-cardiac patients, and antibiotics' side effects were predominantly poor (46.75±14.82, 39.21±33.09 and 20.27±18.77, respectively over 100). Practicing outside Beirut, undergraduate qualification in Lebanon, and post-graduate qualification predicted higher knowledge. 75.9% acknowledged the contribution of dentistry-based prescribing to antibiotic resistance and 94.7% knew at least one cause of resistance.

Conclusions: Dentists show positive attitude towards antimicrobial resistance. Yet, they lack uniformity in antibiotic stewardship. Poor knowledge and guideline-incongruent prophylactic and therapeutic prescribing are observed. Development of targeted interventions is needed to promote judicious antibiotic use within Lebanese dentistry.

Keywords

Antibiotic Prophylaxis; Health Knowledge, Attitudes, Practice; Inappropriate Prescribing; Professional Practice; Guideline Adherence; Penicillins; Streptomycetes; Dentists; Surveys and Questionnaires; Lebanon

INTRODUCTION

Antimicrobial resistance is a serious threat to human life, posing catastrophic public health and economic burdens.¹ Since the mid-1990s, dentistry-based antimicrobial prescribing emerged as one potential driver of the global phenomenon of antibiotic resistance.² Clearly, the use of antibiotics as an adjunct to local treatment is the most appropriate method of managing oral infections.^{3,4} However, its inappropriate prescription would not provide sufficient benefit yet, it runs the risk of causing side effects ranging from gastrointestinal disturbances to fatal anaphylactic shock and emergence of resistant bacteria, and yields greater health.^{5,6} Thus, dentistry-based antibiotic prescribing for prophylactic and therapeutic conditions is dictated by defined criteria, and dentists are urged to judiciously prescribe antibiotics.^{4,7-10} However, the

increasing and inappropriate use of antibiotic by dental professionals remain an international finding.¹¹⁻¹⁷

Knowledge and attitudinal factors are pivotal in explaining this evidence-practice gap.¹⁸ Specifically, in the Middle East, dentists are prone to prescribe on patient's demand, especially when short of time. Antibiotics are abused to prevent postoperative infections or as a consequence of the lack of aseptic clinical techniques.¹⁹ Conflicting data from the region show that in some countries in spite of good knowledge of local and international guidelines, and awareness of the importance of the judicious use of antimicrobials, dentists tend to use antibacterials for inappropriate indications.^{15,20} Studies have shown patterns of overprescribing among dentists where broader spectrum antibiotics, longer durations and higher doses are given.²¹⁻²⁶ In Lebanon, information on antibiotic stewardship in dentistry is scarce. The only available evidence is in acute and chronic dento-alveolar abscess and emanate from a small study conducted in Beirut. It reports results parallel with the international literature: inappropriate use in terms of dosage, duration and frequency is evident, with amoxicillin being the primary prescribed agent.²⁷

Monitoring trends in antibiotic prescriptions by dentists and elucidating pertaining knowledge and attitudinal factors may reveal previously unrecognized opportunities to curb prescribing, and might identify areas of concern in a

Hicham MANSOUR. BDS. Department of Restorative and Esthetic, Faculty of Dentistry, Lebanese University. Beirut (Lebanon). hicham_mansour04@yahoo.com

Mireille FEGHALI. BDS. Department of Restorative and Esthetic, Faculty of Dentistry, Lebanese University. Beirut (Lebanon). mireillefeghali@hotmail.com

Nadine SALEH. PharmD, MS, PhD. Division of epidemiology and biostatistics - Faculty of Public Health - Lebanese University, Beirut (Lebanon). nadeensh@gmail.com

Mona ZEITOUNY. BDS. Department of Restorative and Esthetic, Faculty of Dentistry, Lebanese University. Beirut (Lebanon). monazeitouny@live.com

service or where there is a potential for improvement and optimize antibiotic treatments and stem the emergence and spread of resistance.^{5,18} A national survey was conducted among Lebanese dentists to explore antibiotic use and its concordance with guidelines, and to describe pertaining knowledge and attitudinal factors.

METHODS

An observational cross-sectional telephone-based survey was performed between July and September 2017. The study participants were chosen from the list of Lebanese dentists registered at the Lebanese Order of Dentists. Out of 4432 registered dentists, complete data were obtained for 3222 dentists. Dentists were then sorted according to their region of practice and gathered into subgroups based on the corresponding governorate. They were distributed as follows: 20% from Beirut, 55% from Mount Lebanon, 13% from South Lebanon (including Nabatiyeh), 11% from Bekaa, and 1% from North Lebanon.

The study sample was drawn to respect the same distribution of dentists per governorate. A minimum sample size of 322 participants (10% of the list of dentists with complete data) was considered sufficient to fulfill the study's main objective.

A systematic random sampling was then adopted, and dentists with an odd number in the list {1, 3, 5, 7, etc.} were orderly called until reaching the required number of participants from each region. In total, we had to make 1530 phone calls to be able to reach 460 dentists, among whom 322 gave their oral consent to participate in the study (1070 calls resulted in the following: "dentist absent" or "dentist busy" or "no answer"). The telephonic interview lasted between 10 and 15 minutes.

A standardized questionnaire was designed in English as well as in French. Translations were supervised by professional translators. The questionnaire was pre-tested with 10 dentists for validity and acceptability. Validity was examined by evaluating whether the questions were comprehensive. Acceptability was evaluated by asking the dentists how they found answering the questionnaire and if they wanted to omit or add questions. Confidentiality of the respondent was ensured. The first section of the questionnaire included questions regarding demographic data, specialty, education details, level of experience, working place, attendance of continuing education sessions, average activity. In the second section, dentists were asked to indicate their usual prescription pattern of antibiotics and factors that influence their behavior. The third section was composed of table with a list of different non-invasive and invasive dental procedures and a question about their routine prophylactic or curative prescription of antibiotics (type, dose, duration, route of administration) in general population and in high risk of infection patients (immune-suppressed and with high risk of infective endocarditis). The final section included their knowledge concerning antibiotics, high risky patients, recommendations and their own role in antimicrobial resistance.

The Lebanese University ethics committee waived the need for approval since the study was observational, anonymous and respected the individuals' confidentiality.

Statistical Analysis

Data were collected and all analyses were performed using SPSS version 20. Descriptive analysis was generated. Means and standard deviations were used for quantitative variables while percentages were shown for qualitative variables.

Knowledge questions were isolated and scored. One (1) mark was given for every correct response and zero (0) for an incorrect response. Responses of "Do not know" were counted as incorrect, and no points were given. The total knowledge score was the sum of all correct answers. For dentists who provided answers to all questions, mean knowledge score (%) was calculated and divided into three categories: poor (<60%), intermediate (60-80%) and good (>80%) level.

The antibiotic prescriptions in different dental procedures were compared to recommended guidelines⁵⁻⁹ in order to evaluate their appropriateness (indication, type, dose, frequency and duration). Finally, a bivariate analysis was computed to observe the relations between the knowledge of dentists and their demographic and professional characteristics; i.e. Independent Samples T-Test to explore the association between knowledge scores and independent variables having two mutually exclusive groups, and One-Way ANOVA to explore the association between knowledge scores and independent variables having 3 or more mutually exclusive groups.

RESULTS

322 dentists completed the interview. Their mean age was 44.87 years (9.60; range: 24-67), and 67.1% of them were males. The professional characteristics of participants are provided in Table 1. Reported antibiotic prescribing frequency varied widely among the respondents: on average, 8.8 (11.73) systemic courses were prescribed weekly, and overall 17.51% (18.32%) of dental consultations resulted in the prescription of an antibiotic.

Table 2 details antibiotic prescribing practices. It should also be noted there was a wide range of antibiotics prescribed as a first choice for people who are allergic to penicillin, as well as for both pregnant and lactating women, with varying spectrums of activity. To note that macrolides were the most common first-line antibiotics prescribed to patients allergic to penicillin (47.4%). Interestingly, 5.9% of dentists reported penicillin agents as their first choice. In addition, cetirizine was recommended by one respondent as a first choice antibiotic for a patient allergic to penicillin. Amoxicillin and amoxicillin/clavulanate (Penicillins) were the most common antibiotics prescribed for pregnant and lactating women, followed by macrolides. More than one-quarter of respondents reported referring these women to their gynecologists, when antibiotic prescription is needed (26.9% and 28.5%, respectively). Also, referral of cardiac patients, when necessary, was high (79.4%). 86.9% of the sample always enquired whether their patients are taking antibiotics before proceeding to

Table 1. Professional Characteristics of Participating Dentists						
					N	%
Main Consultation Region (n=322)						
	Mount-Lebanon			177		55.0
	Beirut			65		20.2
	South (including Nabatiyeh)			42		13.1
	Bekaa			35		10.9
	North			3		0.9
Primary dental qualification (n=318)*						
	Lebanon			192		60.4
	Other countries			126		39.6
Years in practice (n=314)*						
	< 1-5 years			21		6.7
	5-10 years			43		13.7
	> 10 years			250		79.6
Specialty (n=309)*						
	General practitioner			134		43.4
	Oral surgeon			33		10.7
	Endodontic			30		9.7
	Implant surgeon			28		9.1
	Pediatric dentist			22		7.1
	Orthodontist			21		6.8
	Restorative Dentist			12		3.9
	Prosthodontics			15		4.9
	Periodontics			9		2.9
	Other			5		1.6
Postgraduate qualification (n=315)*						
	None			134		42.5
	Master's degree			114		36.2
	University Diploma			49		15.6
	PhD			18		5.7
Country where postgraduate qualification was obtained (n=169)*						
	Lebanon			132		78.1
	Western Europe			23		13.6
	Eastern Europe			8		4.7
	USA			5		3.0
	Egypt			1		0.6
Practice setting (n=322)						
	Private Clinic			308		95.7
	Private Hospital			1		0.3
	Public Hospital			1		0.3
	Mixt			12		3.7
Continuing education source (n=314)*						
	None			13		4.1
	National conferences			154		49.0
	National and international conferences and continuing education lectures			99		31.5
	International conferences			33		10.5
	Continuing education lectures			15		4.8
Guidelines followed for prescribing prophylaxis regimens for infective endocarditis among susceptible patients (n=315)*						
	Do not know			103		32.7
	Guidelines provided during dental qualification years			84		26.7
	American Health Association (AHA)			73		23.2
	Agence Française de Sécurité Sanitaire des Produits de Santé (Afsaps)			33		10.5
	National Institute for Clinical Excellence (NICE) and the British Society for Antimicrobial Chemotherapy (BSAC)			21		6.7
	Other			1		0.3
Attending at least 1 lecture relating to the use of antibiotics in dental medicine during the past 5 years (n=316)*					154	48.7
Reading at least 1 journal article relating to the use of antibiotics in dental medicine during the past 5 years (n=314)*					157	50.0
		Min	Max	Median	IQR	Mean
Number of patients per week (n=233)						
		10	240	48.00	30.00	53.71
Number of prescribed systemic antibiotics courses per week (n=260)						
		0	100	5.00	7.00	8.80
Frequency of antibiotic prescription per dental consultation (%) (n=274)						
		0	100	10.00	12.50	17.51
*Valid percentages are reported, Min: minimum; Max: maximum; IQR: interquartile range; SD: standard deviation						

Table 2. Attitude of participating dentists toward antibiotic prescribing		N	%
First choice antibiotic prescribed to patients allergic to penicillin (n=289)*	Spiramycin + Metronidazole	86	29.8
	Spiramycin	69	23.9
	Unspecified Macrolides	35	12.1
	Clindamycin	34	11.8
	Clarithromycin	33	11.4
	Amoxicillin	11	3.8
	Cephalosporin	10	3.5
	Amoxicillin + Clavulanic acid	6	2.1
	Metronidazole	2	0.7
	Cetirizine	1	0.3
	Ciprofloxacin	1	0.3
Sulphamides + Diamonopyrimidine	1	0.3	
First choice antibiotic prescribed to a pregnant woman (n=275)*	Amoxicillin + Clavulanic acid	86	31.3
	Spiramycin	53	19.2
	Amoxicillin	44	16.0
	Spiramycin + Metronidazole	8	2.9
	Clindamycin	4	1.5
	Azithromycin	2	0.7
	Cephalosporin	2	0.7
	Aminosides	1	0.4
	Penicillines or Spiramycin + Metronidazole	1	0.4
	Referral to gynecologist	74	26.9
First choice antibiotic prescribed to a breastfeeding woman (n=274)*	Amoxicillin + Clavulanic acid	85	31.1
	Spiramycin	51	18.6
	Amoxicillin	42	15.3
	Spiramycin + Metronidazole	9	3.3
	Clindamycin	5	1.8
	Cephalosporin	2	0.7
	Gentamycin	1	0.4
	Penicillines or Spiramycin + Metronidazole	1	0.4
	Referral to gynecologist	78	28.5
Frequency of referring cardiac patients to their physician when necessary (n=321)*	Always	255	79.4
	Sometimes	58	18.1
	Never	8	2.5
Enquire if the patient is currently taking an antibiotic before proceeding to consultation (n=320)*	Always	278	86.9
	Often	24	7.5
	Sometimes	14	4.4
	Never	4	1.3
Attitude regarding a patient who has already taken antibiotics before consultation (n=154)*	Continue antibiotic course	85	55.2
	Action depends on the antibiotic	32	20.8
	Change the antibiotic	19	12.3
	Action depends on time (change if antibiotic taken during last month)	8	5.2
	Discontinue antibiotic course	7	4.5
	Continue antibiotic course and add vitamins	2	1.3
	Increase the dose	1	0.6
Feeling pressure from patients to prescribe antibiotics (n=319)*	Always	27	8.5
	Often	43	13.5
	Sometimes	72	22.6
	Never	177	55.0
Factor(s) mostly influencing antibiotics prescribing behavior (n=321)* [†]	Previous antibiotic experience	261	81.3
	Comorbidities of the patient	174	54.2
	Socio-economic status of the patient	103	32.1
	Price of the antibiotic	101	31.5
	Samples availability	44	13.7
	Medical representative visits	37	11.5

*Valid percentages are reported; [†]Percentages may add up to more than 100%, due to multiple possible answers

the consultation. Continuing antibiotic course was the dominant action when a patient was found to be already taking antibiotics (55.2%). Only 5.2% of dentists reported changing the antibiotic if given during the past month. Nearly half (45.0%) of participating dentists reported being, to varying extent, pressured by patients to prescribe antibiotics. Factors governing antibiotic prescribing were primarily physician-related (previous antibiotic experience: 81.3%), followed by patient-related factors (presence of comorbidities: 54.2%). It also should be noted that the socio-economic status (32.1%) and price of the antibiotic (31.5%) were approximately one third of the factors that influenced antibiotic prescribing behavior. Other less influencing factors were the availability of the samples and medical representatives (13.7% and 11.5%, respectively).

Table 3 describes prophylactic antibiotic prescription patterns of sampled dentists. The vast majority of dentists refrained from prescribing antibiotics for restoration

(96.7%), prosthesis (96.4%), crown (93.8%) and local anesthesia (91.6%). Systematic antibiotic prescription was mostly considered for implant (55.7%), bone graft (48.3%) and surgical extraction (mandibular tooth: 46.9%, maxillary tooth: 47.1%). Prescription for patients at high risk for infection was more common for braces (33.3%) and scaling (28.2%). Great divergences were noted for bone graft, implant, teeth extraction and gectomy. Conformity with evidence-practice guidelines was inconsistent; it was high for restoration and interim care (96.7% each), prosthesis (96.4%), crown (93.8%) and local anesthesia (91.6%), where antibiotics are not indicated. Agreement with guidelines was especially low for procedures where prophylactic antibiotics should be prescribed for high-risk patients, such as implant (2.6%), intraligamentary local anesthesia (4.2%), tumor resection (4.6%), frenectomy (8.8%), gingivectomy (9.2%) and Crown lengthening (10.4%). Among those who prescribed prophylactic antibiotics correctly when indicated, conformity with evidence-practice guidelines

Table 3. Prophylactic and Curative Antibiotic Prescribing Practices of Participating Dentists and Conformity with Evidence-Practice Guidelines

N (%)	No	Yes all patients	Yes High-risk patients [‡]	Indication	Conformity with evidence-practice guidelines*		
					Type	Dose	Duration
					Among those who provided a correct answer to indication		
Reported prophylactic antibiotic prescribing*							
Bone graft (n=180)	84 (46.7)	87 (48.3)	9 (5)	87 (48.3)	67 (77.0)	54 (62.1)	3 (3.4)
Braces (n=30)	20 (66.7)	0 (0)	10 (33.3)	20 (66.7)	NA		
Crown (n=306)	287 (93.8)	0 (0)	19 (6.2)	287 (93.8)	NA		
Crown lengthening (n=240)	165 (68.8)	50 (20.8)	25 (10.4)	25 (10.4)	2 (8.0)	1 (4.0)	0 (0)
Extraction mandibular tooth (n=294)	103 (35)	138 (46.9)	53 (18)	138 (46.9)	84 (60.9)	38 (27.5)	4 (2.9)
Extraction maxillary tooth (n=293)	106 (36.2)	138 (47.1)	49 (16.7)	138 (47.1)	91 (65.9)	49 (35.5)	4 (2.9)
Flap surgery (n=166)	101 (60.8)	49 (29.5)	16 (9.6)	101 (60.8)	NA		
Frenectomy (n=249)	188 (75.5)	39 (15.7)	22 (8.8)	22 (8.8)	4 (18.2)	2 (9.1)	1 (4.5)
Germectomy (n=209)	107 (51.2)	80 (38.3)	22 (10.5)	80 (38.3)	51 (63.8)	26 (32.5)	2 (2.5)
Gingivectomy (n=293)	225 (76.8)	41 (14)	27 (9.2)	27 (9.2)	4 (14.8)	2 (7.4)	1 (3.7)
Implant (n=228)	95 (41.7)	127 (55.7)	6 (2.6)	6 (2.6)	2 (33.3)	1 (16.7)	1 (16.7)
Interim care (n=306)	296 (96.7)	0 (0)	10 (3.3)	296 (96.7)	NA		
Intraligamentary local anesthesia (n=311)	298 (95.8)	3 (1)	13 (4.2)	13 (4.2)	2 (15.4)	2 (15.4)	0 (0)
Local anesthesia (n=311)	285 (91.6)	13 (4.2)	13 (4.2)	285 (91.6)	NA		
Necrotic tooth (n=299)	188 (62.9)	84 (28.1)	27 (9)	188 (62.9)	NA		
Prosthesis (n=306)	295 (96.4)	0 (0)	11 (3.6)	296 (96.4)	NA		
Restoration (n=306)	296 (96.7)	0 (0)	10 (3.3)	296 (96.7)	NA		
Scaling (n=309)	215 (69.6)	7 (2.3)	87 (28.2)	215 (69.6)	NA		
<i>Simple extraction (n=305)</i>	209 (68.5)	46 (15.1)	50 (16.4)	50 (16.4)	23 (46.0)	17 (34.0)	1 (2.0)
<i>Tumor resection (n=151)</i>	126 (83.4)	18 (11.9)	7 (4.6)	7 (4.6)	0 (0)	0 (0)	0 (0)
Reported curative antibiotic prescribing*[‡]							
Agressive periodontitis (n=268)	77 (28.7)	176 (65.7)	15 (5.6)	176 (65.7)	90 (51.1)	47 (26.7)	60 (34.1)
Apical abscess (n=306)	97 (31.7)	192 (62.7)	17 (5.6)	17 (5.6)	9 (52.9)	0 (0)	1 (5.9)
Bacterial stomatitis (n=142)	104 (73.2)	38 (26.8)	0 (0)	38 (26.8)	24 (63.2)	21 (55.3)	17 (44.7)
Cellulitis (n=253)	53 (20.9)	174 (68.8)	26 (10.3)	174 (68.8)	122 (70.1)	0 (0)	67 (38.5)
Chronic periodontitis (n=289)	213 (73.7)	55 (19)	21 (7.3)	213 (73.7)	NA		
Combined lesion (n=293)	173 (59)	117 (39.9)	3 (1)	173 (59)	NA		
Fistula (n=285)	137 (48.1)	127 (44.6)	21 (7.4)	127 (44.6)	77 (60.6)	39 (30.7)	58 (45.7)
Gingivitis (n=297)	246 (82.8)	37 (12.5)	14 (4.7)	246 (82.2)	NA		
Maxillary sinusitis (n=159)	104 (65.4)	48 (30.2)	7 (4.4)	48 (30.3)	36 (75.0)	15 (31.3)	18 (37.5)
Osteomyelitis (n=170)	68 (40)	90 (52.9)	12 (7.1)	90 (52.9)	78 (86.7)	61 (67.8)	35 (38.9)
Periapical abscess (n=305)	77 (25.2)	203 (66.6)	25 (8.2)	25 (8.2)	8 (32)	0 (0)	1 (4.0)
Periimplantitis (n=170)	100 (58.8)	51 (30)	19 (11.2)	19 (11.2)	0 (0)	0 (0)	0 (0)
Periodontal abscess (n=284)	75 (26.4)	192 (67.6)	17 (6)	17 (6)	8 (47.1)	4 (23.5)	4 (23.5)
Pulpitis (n=305)	269 (88.2)	27 (8.9)	9 (3)	269 (88.2)	NA		
Salivary gland infection (n=136)	113 (83.1)	23 (16.9)	0 (0)	23 (16.9)	20 (87)	18 (78.3)	9 (39.1)
Tooth decay (n=311)	304 (97.7)	5 (1.6)	2 (0.6)	304 (97.7)	NA		

NA: not applicable. †Dentists describing cases as referred or rarely seen were excluded; *Valid percentages are reported; ‡Selected patients with cardiac conditions; compromised immunity; shunts, indwelling vascular catheters, medical devices; and prosthetic joints (5-9).

types of prophylactic antibiotics		
Amoxicillin		52.5
Amoxicillin and Clavulanic acid		36.4
Unspecified penicillin		6.1
Spyramicin		3.0
Amoxicilin or Spyramicin		1.0
Depends on the case		1.0
doses of prophylactic antibiotics		
2 g		63.9
3 g		9.8
Flash dose		8.2
50 mg/Kg		4.9
1-2 g		3.3
2-3 g		3.3
Other		6.6
timing of antibiotics prophylaxis		
1 hour before procedure		34.0
1 hour before and after procedure		10.6
1 hour before and 6 hours after procedure		7.4
1 hour before and 7 days after procedure		6.4
2 hours before procedure		6.4
1 day before procedure		4.3
2 days before procedure		3.2
3 days before procedure		3.2
Other		24.5

*Valid percentages are reported

regarding the type of antibiotics ranged between 0 and 77%; whereas that of dose ranged between 0 and 62.1%, and that of duration between 0 and 16.7%. Overall, there was a significant divergence from the guidelines for several indications for both patient who were and are not at risk.

Table 3 also shows curative antibiotic prescription patterns of participants; answers were inconsistent for the majority of conditions. Non-prescription was most common in case of tooth decay (97.7%), pulpitis (88.2%), salivary gland infection (83.1%) and gingivitis (82.8%). Around two-thirds

of the dentists reported prescribing antibiotics for all cases diagnosed with cellulitis (68.8%), periodontal abscess (67.7%), periapical abscess (66.6%), aggressive periodontitis (65.7%) and apical abscess (62.7%). Discrepancies were mainly noted for fistula, aggressive periodontitis, apical abscess and maxillary sinusitis. It is important to note that 11.9% of respondents prescribe antibiotics for pulpitis and 17.8% of participants prescribe antibiotics for gingivitis, which is unnecessary prescribing. Also for conditions such as cellulitis (20.9%) and salivary gland infections (83.1%), there were a significant proportion of dentists for both conditions who do not prescribe antibiotics when they are actually indicated. The lowest conformities were observed for apical abscess (5.6%), periodontal abscess (6%) and periapical abscess (8.2%), where curative antibiotics are indicated only for high risk patients. Among dentists who provided a correct answer to indication, the prescribed types of antibiotics were adequate for cases with salivary gland infection (87%) and osteomyelitis (86.7%), and were all inadequate for periimplantitis. When curative antibiotics were prescribed correctly when indicated, conformity with evidence-practice guidelines regarding the type of antibiotics ranged between 0 and 87%; whereas that of dose ranged between 0 and 78.3%, and that of duration between 0 and 45.7%.

As displayed in Table 4, penicillins were the dominant type (95.0%) of prophylactic antibiotics for cardiac patients. Answers were greatly scattered, especially for the dose and timing. Doses ranged between 1.87g up to 5g, with 63.9% prescribing 2g. Timing of antibiotic prophylaxis ranged between 3 days before the procedure, up to 7 days afterwards. The most common timing was 1 hour before procedure (34.0%), followed by 1 hour before and after the procedure (10.6%).

	N	%
<i>Prophylactic prescription of antibiotics for cardiac conditions (correct answers)*</i>		
Prosthetic cardiac valves (n=140)	135	96.4
Rheumatic heart disease (n=131)	26	19.8
Mitral valve prolapsed with valvular regurgitation (n=124)	16	12.9
Previous infective endocarditis (n=127)	102	80.3
Previous coronary artery bypass graft surgery (n=140)	70	50.0
Hypertrophic cardiomyopathy (n=147)	84	26.1
Intravascular cardiac pacemakers (n=147)	49	33.3
Myocardial infarct in the last 6 months (n=125)	26	20.8
Cardiac transplantation recipients who develop cardiac valvulopathy (n=131)	65	49.6
Unrepaired cyanotic heart disease (n=134)	57	42.5
Recently placed coronary stents (n=144)	30	20.8
Atrial septal defect after 6 months of repair (n=134)	59	44.0
Ventricular septal defect with repair (n=140)	56	40.0
Patent ductus arteriosus (n=140)	56	40.0
Cardiac catheterization without stents (less than 1 year) (n=183)	61	43.9
<i>Prophylactic prescription of antibiotics for other conditions (in case of invasive procedure) (correct answers)*</i>		
Human immunodeficiency virus (n=155)	90	58.1
Neutropenia (n=132)	51	38.6
Cancer chemotherapy (n=136)	81	59.6
Diabetes (n=245)	188	76.7
Hematopoietic stem cell or solid organ transplantation (n=133)	56	42.1
Bisphosphonate therapy (n=173)	62	35.8
Chronic steroid usage (n=172)	91	52.9
Asplenism or status post splenectomy (n=175)	69	39.4
Patients with prosthetic joints (n=173)	16	9.2

*Valid percentages are reported

Table 6. Mean Knowledge Scores (%) by Sociodemographic and Professional Characteristics				
		Prophylaxis for cardiac conditions (n=76)	Prophylaxis for non-cardiac conditions (n=85)	Side effects of antibiotics (n=322)
Overall score		46.75 (14.82)	39.21 (33.09)	20.27 (18.77)
Age in years	24-34	53.33 (12.34)	39.35 (35.13)	23.80 (17.77)
	35-50	43.95 (15.27)	43.46 (31.23)	19.23 (18.46)
	>50	45.83 (15.21)	34.78 (33.55)	19.95 (19.95)
	p-value	0.13	0.62	0.30
Gender	Male	46.02 (15.35)	40.50 (33.36)	19.17 (18.65)
	Female	50.00 (12.19)	35.74 (32.81)	22.50 (18.90)
	p-value	0.36	0.55	0.13
Region	Beirut	44.10 (7.47)	23.14 (28.73)	13.40 (10.97)
	Other	47.30 (15.91)	45.53 (32.75)	22.01 (19.92)
	p-value	0.27	0.003	<0.001
Experience years	< 1-5 years	56.19 (13.80)	37.03 (41.94)	25.85 (15.39)
	5-10 years	47.33 (19.98)	40.00 (32.74)	21.26 (18.00)
	> 10 years	45.63 (13.89)	39.65 (31.91)	20.00 (19.28)
	p-value	0.20	0.97	0.38
Specialty	General practitioner	47.13 (15.07)	30.82 (31.65)	20.25 (14.47)
	Other	46.26 (14.71)	44.03 (33.22)	20.28 (21.35)
	p-value	0.80	0.07	0.98
Undergraduate qualification	In Lebanon	52.72 (13.37)	45.89 (31.91)	21.94 (19.87)
	Outside Lebanon	42.17 (14.38)	32.73 (33.43)	18.14 (16.89)
	p-value	0.002	0.07	0.06
Post-graduate qualification	No	47.23 (15.07)	30.82 (31.65)	20.25 (14.47)
	Yes	46.45 (14.90)	46.18 (32.91)	20.59 (21.52)
	p-value	0.84	0.04	0.86
Continuing education	No	46.66 (11.54)	24.07 (32.52)	18.68 (10.72)
	Yes	47.04 (15.26)	43.07 (32.91)	20.45 (19.09)
	p-value	0.96	0.17	0.58
Number of patients per week	0-50	46.23 (18.13)	44.14 (34.19)	21.88 (19.98)
	51-100	49.16 (12.38)	48.41 (31.60)	27.60 (21.85)
	>100	42.85 (14.32)	68.88 (27.66)	19.64 (11.51)
	p-value	0.67	0.29	0.13
Frequency of antibiotic prescription per dental consultation (%)	0-10	45.71 (15.05)	40.54 (34.36)	21.36 (17.17)
	>10	49.85 (16.43)	41.58 (32.71)	21.65 (20.82)
	p-value	0.30	0.89	0.90

Regarding cardiac conditions, the highest knowledge was for prosthetic cardiac valves (96.4%), followed by previous infective endocarditis (80.3%) (Table 5). The adequacy of answers greatly decreased for all other conditions. The worst knowledge was observed for mitral valve prolapsed with valvular regurgitation (12.9%) and rheumatic heart disease (19.8%). The mean score of dentists who provided answers to all questions in this section (n=76) was 46.75 (14.82). None of them had good knowledge about prophylactic prescription of antibiotics for cardiac conditions; two-thirds (67.1%) had poor knowledge, and one-third (32.9%) had intermediate knowledge. Regarding non-cardiac conditions, less than half of respondents could adequately identify prophylactic antibiotic prescription, except for the cases of diabetes (76.7%), cancer chemotherapy (59.6%), infection with the human immunodeficiency virus (58.1%) and chronic steroid usage (52.9%). Knowledge pertaining to prophylactic antibiotic prescription for patients with prosthetic joints was the worst (9.2%). For dentists who provided answers to all

questions in this section (n=85), the average knowledge score was 39.21 (33.09). The participants had predominantly poor knowledge (67.1%); 14.1% had intermediate knowledge and only 18.8% showed good knowledge.

In total, 50.3% of sampled dentists could correctly identify at least one side effect of amoxicillin/co-amoxiclav. This rate sharply declined for other antibiotics, and was almost null for cephalosporin (3.7%). The mean knowledge score about side effects of antibiotics was 20.27 (18.77). Almost all dentists (97.5%) had poor knowledge; only 4 (1.2%) had intermediate knowledge and 4 others (1.2%) exhibited good knowledge.

As shown in Table 6, in the bivariate analysis, demographic and professional characteristics did not influence knowledge scores; with the exception of dentists in Beirut being less knowledgeable of prophylactic prescription for non-cardiac patients and antibiotic side effects than those working in other regions. Moreover, dentists receiving their

undergraduate qualification in Lebanon had greater knowledge scores about prophylactic prescription for cardiac patients than the others, and those with a post-graduate qualification had higher knowledge of prophylactic prescription for non-cardiac patients than their peers.

Finally, 75.9% of respondents were aware of the contribution of dentistry-based antibiotic prescribing to the problem of antibiotic resistance at the national level and 94.7% knew at least one cause of antibiotic resistance.

DISCUSSION

Given the potential contribution of dentistry-based antibiotic misuse to the epidemic of antimicrobial resistance, this study was the first effort to describe current knowledge, attitude and practices related to antibiotics, and to assess the extent to which prophylactic and therapeutic prescribing conforms to guidelines among dentists across Lebanon. In order to reach the desired sample size (322 dentists), the survey targeted 1,530 dentists, of whom, 460 were accessible, revealing a participation rate of 21% out of all targeted dentists, and a response rate of 70% among those who were accessible. This is in line with previous similar national studies conducted among dentists in other countries.^{13,17,28}

Although, within the population studied, the reported rate of antibiotic prescribing was relatively high (17.51%) compared to other studies in Australia, Belgium and the United Kingdom^{12,29,30}; and while participants lacked uniformity in antibiotic prescribing knowledge and practices, unindicated, inappropriate and extended uses were obvious, suggesting guideline-incongruent prophylactic and therapeutic prescribing. The problematic prescribing in Lebanon is further evidence to the international concern of dentistry-based antibiotic misuse^{11-17,28}, and provides additional argumentation justifying the solicitation of national efforts to promote judicious antibiotic use across the profession. Several factors noted in our sample emerge as potential contributors to these findings, including poor knowledge of evidence-practice regimens, limited exposure to scientific updates relating to the use of antibiotics, in addition to pressure of non-medical factors, such as patient requests for antibiotics prescription and influence of pharmaceutical industry. Various non-clinical pressures are in agreement with studies from other countries.^{11,12,31,32} Our sample exhibited several conform prescribing behaviors, such as mainly using macrolides as first-line antibiotics for patients allergic to penicillins.³³ This behavior was in line with data from Belgium¹², yet differed from data reported from other countries, where clindamycin and erythromycin were the most prescribed antibiotics in the United Arab Emirates and United Kingdom^{13,15} and in Iran¹⁶, respectively. Several factors might explain this finding, among them is the comparative safety and tolerance of macrolides and the concern from the higher rates of fatal and nonfatal adverse drug reactions associated with *C. difficile* infections with clindamycin use³⁴ in one hand, and the unavailability of erythromycin in oral form in Lebanon, on the other hand. Yet, several deviant practices related to this condition were observed, such as the use of penicillins for these patients,

or even substituting antibiotics by anti-histaminic or even not recognizing that amoxicillin and amoxicillin/clavulanic acid are both from the penicillin group. These behaviors-denoting poor knowledge of basic antibiotic pharmacology-might engender serious side effects, some of which could be life-threatening. Similarly, as previously noted in Lebanon³⁵, the use of penicillins as primary antibiotics for pregnant and lactating women was evident. Yet, a substantial proportion of dentists adopted metronidazole as their first choice for these women. First-line use of this agent is not supported by evidence, especially during the first semester of gestation and during lactation, rather, it is typically indicated for second-line use.³⁴ Although few in numbers, alarming practices emerged in this patient population, such as the use of spiramycin, aminosides and gentamycin in first-line.

On the other hand, our sample showed evidence of factors fostering antimicrobial resistance. First, antibiotic prescribing was found to be biased toward broad spectrum agents, i.e. association amoxicillin with clavulanic acid, association spiramycin with metronidazole and metronidazole, which were used in numerous instances, even when not clinically required. This finding is universal among dental practitioners.^{12,14,15,17,20} Second, a high proportion of dentists inquired whether the patient is using antibiotics before consultation; however the vast majority resorted to systematically changing the antibiotic to combat a potential or present infection and only few patients followed the recommendation of changing the antibiotic if taken in the previous month.³⁴ Third, massive doses ranging up to 5g and long duration extending to 8 days of prophylaxis were prescribed for cardiac patients-clearly exceeding the recommended dose and duration of use.³⁴ Fourth, a considerable proportion of physicians adopted routine prescription to all patients, even when not indicated, such as in flap surgery, implant and necrotic tooth; and this misuse was accentuated in antibiotic therapy, such as with cases diagnosed with combined lesion, periapical and periodontal abscess and periimplantitis. Additionally, among physicians who practiced indicated prescribing, optimal adherence to guidelines (type, dose and duration of antibiotic use) was practically inexistent in prophylaxis; it was slightly better for therapeutic use. It is worthy to note that most deficiencies revolved around over and extended use, rather than the type of antibiotics. Finally, the lowest conformity to guidelines was found where antibiotics are indicated for high risk patients only. Potentially, the practitioners might not be confident in identifying high risk patients requiring antibiotics, and resorted the routine prescribing as a preventive mechanism.

As found in other countries^{15,16,35}, knowledge related to conditions where prophylaxis is indicated varied widely amongst participating dentists, was on average far from being optimal, and showed to be specifically low when it comes to non-cardiac conditions. The high referral rate witnessed in our sample, might partially contribute to this finding in a vicious circle. Potentially, dentists are deferring providing care to at-risk patients due to deficiency in their medical knowledge - as noted among other physicians³⁶, thus losing motivation to continuously upgrade their knowledge and skills to take in charge these patients. In

parallel, knowledge about the side effects of antibiotic showed to be the poorest. Addressing this issue is of utmost importance, taking into account the fatal and nonfatal adverse reactions associated antibiotic use. We were able to identify few inconsistent factors associated with higher knowledge: practicing outside Beirut area, receiving their undergraduate qualification in Lebanon, and having a post-graduate qualification.

This study raises many questions to be explored in future endeavors. First, as found in previous publications^{15,18,20}, the majority of dentists were aware of the contribution of dentistry-based antibiotic prescribing to the problem of antibiotic resistance at the national level, and the vast majority of them acknowledged either over, extended and/or misuse of antibiotics as causes of antimicrobial resistance. It was noted that in our sample more importance was accorded to preventing and treating infections rather than preventing antimicrobial resistance. In fact, qualitative data from the United Kingdom indicate that while dentists are aware of the theoretical contribution of dentistry-based prescribing to the emergence of resistance, they perceive it to be far less incriminated than the contribution of their medical col-leagues.¹⁸ This might partly explain the conflicting results emanating from our study. Second, our sample exhibited high referral of pregnant and lactating women, as well as cardiac patients to specialist physicians, when antibiotic prescription is needed. This behavior possibly denotes the limited knowledge, capacity or time of participants to take in charge these critical conditions, or could be regarded as part of the multidisciplinary approach to patient care. Available data do not permit us to generate a conclusion. The study relied on self-reported practices and the answers were not verified against patient records. Participants might have provided more professionally desirable answers, probably resulting in an underestimation of the true prescribing levels. Future studies should consider auditing patient records to provide documented data and ensure accuracy. Another limitation

of this study is the absence of published national treatment guidelines of antibiotics prescription in dental practice and the use of international guidelines to assess conformity which may have created some underestimation of the conformity. Moreover, telephone interviews may have underestimated the real percentage of antibiotic prescription. Finally, we used a systematic random sample which also limits the selection bias. In spite of this, the low response rate may affect the external validity.

CONCLUSIONS

To conclude, while this study pioneers in revealing antibiotic-related knowledge, attitude and practices of dentists in Lebanon, following studies must further investigate the determinants of poor knowledge, attitudinal barriers and inappropriate prescribing, and future research is therefore required to identify practitioners most at-risk of prescribing antibiotics when they are unlikely to be of clinical benefit.

It is now vital that Lebanese professional dental bodies strengthen the knowledge of dentists, and support and encourage judicious antibiotic prophylactic and therapeutic antibiotic prescribing across the profession. Effective interventions could use pharmacist-delivered academic detailing³⁷ as well as clinical audit³⁸ with the issuing of national guidelines and an educational component³⁹, among others.

CONFLICT OF INTEREST

All authors declare no conflict of interest.

FUNDING

This research did not receive any specific grant from funding or sponsorship.

References

1. World Health Organization. The evolving threat of antimicrobial resistance: options for action. Geneva: WHO; 2012.
2. Sweeney LC. Antibiotic resistance in general dental practice--a cause for concern? *J Antimicrob Chemother*. 2004;53(4):567-576. doi: [10.1093/jac/dkh137](https://doi.org/10.1093/jac/dkh137)
3. Fouad AF, Byrne BE, Diogenes AR, Sedgley CM, Cha BY. AAE Guidance on the use of Systemic Antibiotics in Endodontics, AAE Position Statement. 2017. Available at: https://www.aae.org/specialty/wp-content/uploads/sites/2/2017/06/aae_systemic-antibiotics.pdf (accessed May 23, 2018).
4. Guidance Development Group. Drug Prescribing for Dentistry: Scottish Dental Clinical Effectiveness Programme; 2016. Available at: <http://www.sdcep.org.uk/wp-content/uploads/2016/03/SDCEP-Drug-Prescribing-for-Dentistry-3rd-edition.pdf> (accessed May 23, 2018).
5. Al-Haroni M. Bacterial resistance and the dental professionals' role to halt the problem. *J Dent*. 2008;36(2):95-103. doi: [10.1016/j.jdent.2007.11.007](https://doi.org/10.1016/j.jdent.2007.11.007)
6. Lockhart PB, Blizzard J, Maslow AL, Brennan MT, Sasser H, Carew J. Drug cost implications for antibiotic prophylaxis for dental procedures. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2013;115(3):345-353. doi: [10.1016/j.oooo.2012.10.008](https://doi.org/10.1016/j.oooo.2012.10.008)
7. Agence Française de Sécurité Sanitaire des Produits de Santé. Prescription des antibiotiques en pratique bucco-dentaire. *Recommandations de bonne pratique*. AFSSAPS; 2011. Available at: https://www.ansm.sante.fr/content/download/5297/52416/version/12/file/Reco-Prescription-des-antibiotiques-en-pratique-buccodentaire_Septembre2011.pdf (accessed May 23, 2018).
8. American Academy on Pediatric Dentistry Clinical Affairs Committee, American Academy on Pediatric Dentistry Council on Clinical Affairs. Guideline on antibiotic prophylaxis for dental patients at risk for infection. *Pediatr Dent*. 2008-2009;30(7 Suppl):215-218.
9. Wilson W, Taubert KA, Gewitz M, Lockhart PB, Baddour LM, Levison M, Bolger A, Cabell CH, Takahashi M, Baltimore RS, Newburger JW, Strom BL, Tani LY, Gerber M, Bonow RO, Pallasch T, Shulman ST, Rowley AH, Burns JC, Ferrieri P,

- Gardner T, Goff D, Durack DT; American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee; American Heart Association Council on Cardiovascular Disease in the Young; American Heart Association Council on Clinical Cardiology; American Heart Association Council on Cardiovascular Surgery and Anesthesia; Quality of Care and Outcomes Research Interdisciplinary Working Group. Prevention of infective endocarditis: Guidelines from the American Heart Association: A Guideline from the American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee, Council on Cardiovascular Disease in the Young, and the Council on Clinical Cardiology, Council on Cardiovascular Surgery and Anesthesia, and the Quality of Care and Outcomes Research Interdisciplinary Working Group. *Circulation*. 2007;116(15):1736-1754. doi: [10.1161/CIRCULATIONAHA.106.183095](https://doi.org/10.1161/CIRCULATIONAHA.106.183095)
10. Sollecito TP, Abt E, Lockhart PB, Truelove E, Paumier TM, Tracy SL, Tampi M, Beltrán-Aguilar ED, Frantsve-Hawley J. The use of prophylactic antibiotics prior to dental procedures in patients with prosthetic joints. *J Am Dent Assoc*. 2015;146(1):11-16. doi: [10.1016/j.adaj.2014.11.012](https://doi.org/10.1016/j.adaj.2014.11.012)
 11. Marra F, George D, Chong M, Sutherland S, Patrick DM. Antibiotic prescribing by dentists has increased – Why?. *J Am Dent Assoc*. 2016;147(5):320-327. doi: [10.1016/j.adaj.2015.12.014](https://doi.org/10.1016/j.adaj.2015.12.014)
 12. Mainjot A, D'Hoore W, Vanheusden A, Van Nieuwenhuysen JP. Antibiotic prescribing in dental practice in Belgium: Antibiotic prescribing. *Int Endod J*. 2009;42(12):1112-1117. doi: [10.1111/j.1365-2591.2009.01642.x](https://doi.org/10.1111/j.1365-2591.2009.01642.x)
 13. Ireland RS, Palmer NO, Lindenmeyer A, Mills N. An investigation of antibiotic prophylaxis in implant practice in the UK. *Br Dent J*. 2012;213(8):E14. doi: [10.1038/sj.bdj.2012.960](https://doi.org/10.1038/sj.bdj.2012.960)
 14. Cope AL, Francis NA, Wood F, Chestnutt IG. Antibiotic prescribing in UK general dental practice: a cross-sectional study. *Community Dent Oral Epidemiol*. 2016;44(2):145-153. doi: [10.1111/cdoe.12199](https://doi.org/10.1111/cdoe.12199)
 15. Alkhabuli J, Kowash M, Shah A. Knowledge and Attitude of Northern Emirates Dental Practitioners towards Antibiotic Prescription and its Resistance. *Int J Dent Oral Health*. 2016;2:177. doi: [10.16966/2378-7090.177](https://doi.org/10.16966/2378-7090.177)
 16. Kakoei S, Raoof M, Baghaei F, Adhami S. Pattern of antibiotic prescription among dentists in Iran. *Iran Endod J*. 2007;2(1):19-23.
 17. Falkenstein S, Stein JM, Henne K, Conrads G. Trends in antibiotic use and microbial diagnostics in periodontal treatment: comparing surveys of German dentists in a ten-year period. *Clin Oral Investig*. 2016;20(8):2203-2210. doi: [10.1007/s00784-016-1722-6](https://doi.org/10.1007/s00784-016-1722-6)
 18. Cope AL, Wood F, Francis NA, Chestnutt IG. General dental practitioners' perceptions of antimicrobial use and resistance: a qualitative interview study. *Br Dent J*. 2014;217(5):E9. doi: [10.1038/sj.bdj.2014.761](https://doi.org/10.1038/sj.bdj.2014.761)
 19. Dar-Odeh NS, Abu-Hammad OA, Al-Omiri MK, Khraisat AS, Shehabi AA. Antibiotic prescribing practices by dentists: a review. *Ther Clin Risk Manag*. 2010;6:301-306. doi: [10.2147/TCRM.S9736](https://doi.org/10.2147/TCRM.S9736)
 20. Al-Harthi SE, Khan LM, Abed HH, Alkreaty HM, Ali AS. Appraisal of antimicrobial prescribing practices of governmental and non-governmental dentists for hospitals in the western region of Saudi Arabia. *Saudi Med J*. 2013;34(12):1262-1269.
 21. Salako NO, Rotimi VO, Adib SM, Al-Mutawa S. Pattern of antibiotic prescription in the management of oral diseases among dentists in Kuwait. *J Dent*. 2004;32(7):503-509. doi: [10.1016/j.jdent.2004.04.001](https://doi.org/10.1016/j.jdent.2004.04.001)
 22. Alattas HA, Alyami SH. Prescription of antibiotics for pulpal and periapical pathology among dentists in southern Saudi Arabia. *J Glob Antimicrob Resist*. 2017;9:82-84. doi: [10.1016/j.jgar.2017.01.012](https://doi.org/10.1016/j.jgar.2017.01.012)
 23. Vessal G, Khabiri A, Mirkhani H, Cookson BD, Askarian M. Study of antibiotic prescribing among dental practitioners in Shiraz, Islamic Republic of Iran. *East Mediterr Health J*. 2011;17(10):763-769.
 24. Lisboa SM, Martins MA, Castilho LS, Souza e Silva ME, Abreu MH. Prescribing errors in antibiotic prophylaxis by dentists in a large Brazilian city. *Am J Infect Control*. 2015;43(7):767-768. doi: [10.1016/j.ajic.2015.03.028](https://doi.org/10.1016/j.ajic.2015.03.028)
 25. Ryalat S, Hassona Y, Al-Shayyab M, Abo-Ghosh M, Sawair F. Dentists' knowledge and practice regarding prevention of infective endocarditis. *Eur J Dent*. 2016;10(4):480-485. doi: [10.4103/1305-7456.195158](https://doi.org/10.4103/1305-7456.195158)
 26. Koyuncuoglu CZ, Aydin M, Kirmizi NI, Aydin V, Aksoy M, Isli F, Akici A. Rational use of medicine in dentistry: do dentists prescribe antibiotics in appropriate indications? *Eur J Clin Pharmacol*. 2017;73(8):1027-1032. doi: [10.1007/s00228-017-2258-7](https://doi.org/10.1007/s00228-017-2258-7)
 27. Asmar G, Cochelard D, Mokhbat J, Lemdani M, Haddadi A, Ayoubz F. Prophylactic and therapeutic antibiotic patterns of lebanese dentists for the management of dentoalveolar abscesses. *J Contemp Dent Pract*. 2016;17(6):425-433.
 28. Halboub E, Alzaili A, Ali Quadri MF, Al-Haroni M, Al-Obaida MI, Al-hebshi NN. Antibiotic prescription knowledge of dentists in Kingdom of Saudi Arabia: An online, country-wide survey. *J Contemp Dent Pract*. 2016;17(3):198-204.
 29. Teoh L, Stewart K, Marino RJ, McCullough MJ. Part 1. Current prescribing trends of antibiotics by dentists in Australia from 2012 to 2016. *Aust Dent J*. 2018 [ahead of print]. doi: [10.1111/adj.12622](https://doi.org/10.1111/adj.12622)
 30. Lewis MA. Why we must reduce dental prescription of antibiotics: European Union Antibiotic Awareness Day. *Br Dent J*. 2008;205(10):537-538. doi: [10.1038/sj.bdj.2008.984](https://doi.org/10.1038/sj.bdj.2008.984)
 31. Palmer NA, Pealing R, Ireland RS, Martin MV. A study of therapeutic antibiotic prescribing in National Health Service general dental practice in England. *Br Dent J*. 2000;188(10):554-558.
 32. Germack M, Sedgley CM, Sabbah W, Whitten B. Antibiotic Use in 2016 by Members of the American Association of Endodontists: Report of a National Survey. *J Endod*. 2017;43(10):1615-1622. doi: [10.1016/j.joen.2017.05.009](https://doi.org/10.1016/j.joen.2017.05.009)
 33. Brunton L, Hilal-Dandan R, Goodman LS. Goodman and Gilman manual of pharmacology and therapeutics. New York: Mcgraw Hill Professional; 2013.
 34. Thornhill MH, Dayer MJ, Prendergast B, Baddour LM, Jones S, Lockhart PB. Incidence and nature of adverse reactions to antibiotics used as endocarditis prophylaxis. *J Antimicrob Chemother*. 2015;70(8):2382-2388. doi: [10.1093/jac/dkv115](https://doi.org/10.1093/jac/dkv115)
 35. Halboub E, Alzaili A, Ali Quadri MF, Al-Haroni M, Al-Obaida MI, Al-hebshi NN. Antibiotic prescription knowledge of dentists in Kingdom of Saudi Arabia: An online, country-wide survey. *J Contemp Dent Pract*. 2016;17(3):198-204.
 36. Ringberg U, Fleten N, Forde OH. Examining the variation in GPs' referral practice: a cross-sectional study of GPs' reasons for referral. *Br J Gen Pract*. 2014;64(624):e426-e433. doi: [10.3399/bjgp14X680521](https://doi.org/10.3399/bjgp14X680521)

37. Seager JM, Howell-Jones RS, Dunstan FD, Lewis MAO, Richmond S, Thomas DW. A randomised controlled trial of clinical outreach education to rationalise antibiotic prescribing for acute dental pain in the primary care setting. *Br Dent J.* 2006 Aug 26;201(4):217-222. doi: [10.1038/sj.bdj.4813879](https://doi.org/10.1038/sj.bdj.4813879)
38. Zahabiyoun S, Sahabi M, Kharazi MJ. Improving knowledge of general dental practitioners on antibiotic prescribing by raising awareness of the Faculty of General Dental Practice (UK) Guidelines. *J Dent (Tehran).* 2015;12(3):171-176.
39. Palmer NA, Dailey YM, Martin MV. Pharmacology: Can audit improve antibiotic prescribing in general dental practice? *Br Dent J.* 2001;191(5):253-255. doi: [10.1038/sj.bdj.4801156a](https://doi.org/10.1038/sj.bdj.4801156a)